

BOSTON PUBLIC LIBRARY



3 9999 10255 791 3

Accessions

201353

3847.49
Shelf No.

~~#58.28~~



Received Aug. 3, 1876

S

[A

B
PLAT

L

28

1



Buttercup

2



Traveler's Joy

3



Wood Anemone

4



Flanders Poppy

5



Globe Flower

6



Columbine

7



Monk's Hood

8



White Water Lily

9



Prickly Poppy

10



Yellow Horn Poppy

11



Dyer's Woad

12



Whitlow Grass

WILD FLOWERS WORTH NOTICE

A

Selection of some of our Native Plants,

WHICH ARE MOST ATTRACTIVE FROM

THEIR BEAUTY, USES, OR ASSOCIATIONS.

BY

MRS. LANKESTER,

AUTHOR OF 'A PLAIN AND EASY ACCOUNT OF BRITISH FERNS.'

Fully Illustrated by J. E. Sowerby.

LONDON:

HARDWICKE AND BOGUE,

192 PICCADILLY.

Other edition of 3847-72

201,353

Aug. 3, 1846

1158.28

PREFACE.

MY little book on Ferns has been so kindly and well received, that I have been encouraged to supplement it with these pages on "Wild Flowers worth Notice." It has been a difficult task to make a selection,—for what flowers are not worth notice? As, however, this cannot pretend to be an exhaustive treatise on the British Flora, such as exists in many forms and in large ponderous volumes compiled by profound and learned botanists, I have endeavoured to choose such plants as are representatives of particular families, and are remarkable either for their beauty of appearance or useful properties, and to give the best botanical description I can either find or make of them, so as to insure their recognition with the aid of the plate, and to add such traditions, legends, and poetical fancies, as are associated with them, in order to increase the interest with which they may be regarded.

I lately heard a very eloquent and popular preacher discourse of wild flowers and the "lilies of the field," in a strain so completely in harmony with my own thoughts while I had been busy during the week in writing of all their charming and modest beauty, that I felt he too must be a lover of wild flowers and have himself rejoiced in these blessed gifts of beauty which gladden the earth, and remove from man one of the great evils of his fall. Surely, said the preacher, we may regard all that is lovely in nature, the trees, grass, blue sky, sunshine, and above all, wild flowers, as secondary blessings, which we receive and enjoy by virtue of the covenant made with Adam. No thanks to the husbandman that they spring up on every side, gladdening our hearts and cheering our lives; there they are each in its season, without care, without tending. The Great Architect of this glorious world depends not on his creatures for the preservation of these adornments of his universe, but has implanted within each tiny cup or bell that which shall perpetuate itself, and from age to age shall gladden the heart of the poorest wayfarer. "Consider the lilies of the field, how they grow; they toil not, neither do they spin: and yet I

say unto you that even Solomon in all his glory was not arrayed like one of these."

Longfellow well says—

"Wond'rous truths, and manifold as wond'rous,
God hath written in the stars above;
But not less in the bright flow'rets under us
Stands the revelation of His love."

Then the natural connection between wild flowers and bright sunshine, or the first warm days of spring, does it not recall many a pleasant ramble to those who are in the enjoyment of youth and health? And even to the feeble or afflicted, the remembrance of the soft lulling influence of a summer's day, in sweet rural scenes, when everything seems joyous and yet tranquil, is a refreshment and a delight. In a charming series of short essays, called "The Recreations of a Country Parson," is one "Concerning Summer Days," which is so full of pleasant thoughts and the love of green trees and fields, hedges and hedge-rows, that I cannot but wish he would also write "Concerning Wild Flowers." When I first began to write of "Wild Flowers," it was suggested to me that I should select only those susceptible of cultivation; but to me, the great charm of the whole subject is to fancy the beautiful creatures in their natural homes, where they love to grow, not where they are artificially placed and tended by the hand of man. The wild bryony and clematis climbing luxuriantly over hedge and neighbouring tree—none the less rich for the demands made upon it by groups of happy smiling children for wreaths and festoons of wonderful length. The blue forget-me-not peeping out from its bed of green leaves by the side of the clear running stream, asking only for moisture to fringe its sides with turquoise flowers. A hundred other lovely "children of the earth," as the blind girl of Bulwer calls them, owe much of their charm to the "lap" from which they spring—fresh and untouched by the hand of man. Not that I would in any manner depreciate the gardener's art or the skill of the florist, in so tending and cultivating even our native plants, as to produce such perfection of colour and symmetry of form, that it is difficult to recognize our friends of the wayside in the beauties of the garden. But this is surely the admiration with which we regard the well-dressed and fashionable denizens of a city in contrast with the more simple, but, perhaps, not less refined rustic beauties. Then these favourites of ours must be sought for,—they call forth the energy and self-denial of their admirers, and while making great demands in the shape of country walks, and mountain rambles, bestow on their captors rosy cheeks, the inestimable prize of healthful and vigorous frames.

I have often, when urging the necessity of long walks and frequent exercise, been told by young folks living in the midst of rustic lanes, "There is no object to go out for. In cities and towns there are a hundred objects, and we are thus beguiled into walking." Why not then secure an object, if but one, for a country walk; an object which will unfailingly repay you and be cheering in the remembrance? Cultivate an acquaintance with the wild flowers of your own district, study them, gather them, transplant them if you will into a corner of your own garden; but, above all, visit them in their own homes, and be not satisfied till you have made a tolerable friendship with most of our British plants. Like all things of beauty, they perish quickly; and though each month of the year brings its own attractions with it, from the snowdrop of the early spring to the misletoe and lichen of dark December, the lover of flowers will like to preserve the forms of as many favourites as possible, by drying; and few who have botanized in youth, cannot moralize in maturer age over the remembrances thus furnished, and few who have once engaged in collecting plants ever lose the recollection of the study or the interest it inspired. I therefore say to all, observe, collect, and preserve, the wild flowers you find; arrange and name them scientifically, if possible; but if that be not in your power, still they will always be interesting to you as a pleasing record of "times and places, and old familiar faces," which one day you will value. When you have become well acquainted with the wild plants of your own neighbourhood, there are always *rare ones* to look for, and great is the interest and excitement attending the discovery of a rare species in an unexpected locality; but before you can expect to become a discoverer, you must be a tolerable botanist. But botany, like other sciences, requires earnest and systematic study. Those who wish to be able to discover the name of a plant by the aid of botanical books, must first thoroughly understand the structure of a plant. Such a knowledge may be gained by the study of the late Professor Henfrey's "Rudiments of Botany," or Dr. Dresser's "Popular Manual of Botany." When the structure has been mastered, then the descriptions in such volumes as Babington's "Manual of Botany," and Bentham's "British Flora;" or the larger works of Smith, Withering, and Sowerby, may be used with success.

Let it not be supposed, however, that this search after wild flowers need be confined to such as have at their disposal all the appliances of science. The poorest inhabitant of a cottage has within her reach the same delight from this pursuit as the lady of the mansion, and we have many instances of the successful cultivation of botany by those who have to labour hard for their daily bread. Among the hills which surround the great

manufacturing city of the North, Manchester, and even within the very atmosphere of its smoke, there exists, at this day, a club of working naturalists—chiefly botanists. All of them are artisans in some one of the great factories of the district. An account published by one of themselves of their weekly botanical excursions, their field-days, and the healthful and exhilarating effect on the minds and bodies of the members of this club, is most encouraging and delightful. The actual longevity of these humble naturalists is very remarkable. Old Crowther, one of the earliest workers in this direction, died at the age of 79; he was a simple-hearted man, willing to travel any distance and undergo any fatigue, so that he secured his flower. As one of his old companions said, “he was not *learned*, but he was very *loving*.” He never touched his wages for the purpose of botanical pleasure, but took home every penny, and trusted to fortunate accidents for the means of supplying his scientific wants. An account of the life and labours of another of this noble fraternity, the late Samuel Gibson, of Hebden Bridge, appeared in the *Manchester Guardian*, of the 30th May, 1849. His herbarium of plants was sold after his death for the sum of £75, and many portions of his collection are now to be seen in the Peel Park Museum, Salford. In 1858, an annual meeting of these working men naturalists took place near Manchester, at which there mustered not less than 200 zealous and well-informed botanists, all, with the exception of four or five, of the artisan class. The one striking feature of this meeting seems to have been the hale and hearty appearance of the men already advanced in life; they were fine specimens of youth carried on into old age. There is evidently something in natural history wonderfully promotive of length of days. Men never step into the presence of nature with affection and reverence, but they come back blessed and strengthened with a reward.

We have a noble example of what may be done in this direction, even at an early age, in the labours of the late Rev. Professor Henslow, of Hitcham, in Suffolk. The interest he took in his parish schools induced him to think that the girls might with advantage know something of the plants and vegetation of their own village. With his characteristic energy and goodness, he set about teaching all who chose to learn, the elements of botany, and by the encouragement of his own kindly smile and approving words, he soon succeeded in establishing a genuine love for plants, not only to *look at*, but to *understand*, in the minds of these children. The good professor arranged for his pupils a system of naming, classifying, and drying their specimens, and but few girls in the village of Hitcham are now unacquainted with their native plants. A good collection, made, dried, and named by these young botanists, may be inspected any day in the educational

department of the South Kensington Museum, where it is deposited as an example and an encouragement to other village schools. It is but right to add that the testimony of the Inspectors of Schools goes to prove that this school ranks far above the average of schools in the district in every respect; and that in no way is there any difference in its rules or arrangements, with the exception of the introduction of botany. We can, perhaps, readily account for the indirect influence of this study on the habits and minds of the pupils. The attention it awakens, the methods of arrangement and order it encourages, and the accuracy it necessitates, must re-act, in a great measure, on the whole character and thoughts of the learner.

I am glad here to refer to the encouragement lately given to scientific instruction in schools by the Committee of Council on Education; and it would be well that not only our national and village school-teachers should avail themselves of the advantages thus offered, but that our middle and higher-class schools should follow the example. At present the movement is in its infancy, and the full bearing and tendency of offering certificates and prizes to authorized teachers is scarcely understood; but, from the working of the matter since its first introduction, we may fairly predict that unless our sons and daughters awake to a sense of the necessity for exertion and study in this direction, they will be left far behind in the knowledge of the world they live in by the boys and girls of village schools. A series of lectures has been published by the Examiners in Science in the Department of Education, "addressed to teachers on preparation for obtaining science certificates, and the method of teaching a science class." To the one on Botany, price 2*d.*, I would specially refer as explaining and amplifying much that I should wish to say here on the nature and advantages of this study. A Directory, price 6*d.*, is also published, containing minute particulars as to regulations for establishing and conducting science classes in schools. Either of these pamphlets may be obtained by application to the Secretary of the Science and Art Department, South Kensington Museum. The late Rev. Professor Henslow, ever anxious to assist and gratify others at any personal sacrifice, sent me, in spite of the severe and painful illness which carried him off, a little pamphlet prepared as a companion to the specimens of his school plants deposited in the Kensington Museum, entitled "Illustrations to be employed in Practical Lessons on Botany." In it he gives full particulars of the system he has pursued in his schools at Hitcham, and every direction for the commencement of the study of Botany, with little wood-cuts as unmistakable guides. Nearly every book on botany contains directions as to drying plants, and in my "Plain and Easy Account of British Ferns," I have endeavoured to be as clear as

possible on this point. Here, therefore, I would only say to those who wish to make a collection of dried plants, get Bentall's drying-paper, and two boards strapped together with two strong leather straps sufficient to produce considerable pressure. Change your paper often while your plants are drying; when dry, put them down carefully with bits of gummed paper on foolscap sheets; write the name, order, locality and date of finding, neatly underneath. It is a good plan to have a sheet of thick cartridge or brown paper for each family, to enclose all the specimens belonging to that family. You can then place these cases on shelves, in drawers, or a portfolio, for safety and preservation.

Before closing these remarks I would refer to Mr. J. E. Sowerby's "British Wild Flowers," as the only volume with which I am acquainted, containing in a small compass drawings of every known British plant, so carefully executed as to require little or no explanation for the purpose of identifying any special plant by its aid. It seemed to me, however, that it would neither be inappropriate nor unprofitable to show in a selected number of specimens the sort of interest which attaches to every one of these plants, and in what way and with how much enjoyment their acquaintance may be cultivated and improved.

"God made the flowers to beautify
The earth, and cheer man's careful mood;
And he is happiest who has power
To gather wisdom from a flower,
And wake his heart in every hour
To pleasant gratitude." WORDSWORTH.

INTRODUCTION.

THIS little volume certainly does not presume to be a work on Botany, strictly so called, of which there are so many and such exhaustive guide-books to the British Flora. It is not my intention to describe each Natural Order or Family to which our specimens belong. Nearly every reader who is sufficiently interested in plants to care to look for them, and to compare them with the plates and descriptions here given, will already know what constitutes a family, or natural order; but lest some one should take up this book who has not given any previous attention to the subject, I would say that the whole vegetable kingdom is divided into three great classes. These, again, are subdivided into orders, families, or tribes, according as they most resemble each other; these are again subdivided into genera, and again into species. We may popularly explain the system of classifying, thus:—In a library there shall be a number of volumes of all sizes, shapes, and containing varieties of matter; we agree to classify them not only by size and colour, but according to their contents. Take all those treating of chemistry, then all those on botany; let those be families or orders; then all the red-bound chemical books,—that is a genus; then from those separate the volumes with gilt leaves of a certain size, and those with marbled leaves, of a certain size,—these constitute species. In this way have our great botanists divided the vegetable kingdom for the convenience of arrangement and study; and although there are often differences of

opinion as to what shall constitute a *species* or a *variety*, which would correspond in a library to a red, gilt-edged book, with a slight change of style in the cover, say with a thicker, stouter back than the rest; yet in the main there is but little difficulty in preserving order and in classifying every known plant according to its peculiarities. It is very necessary for any one who wishes to study Botany to understand thoroughly the distinguishing points of each natural order. When once these are fixed in the mind, it is easy to recognize plants as belonging to certain families, and from the established habits of the family to draw conclusions as to the nature, locality, and properties of the specimen under notice. In order to give some notion of the nature of a Natural Order or Family, I think it well to give a sketch of that to which our first specimen belongs, and which comes first in nearly all works on Botany.

RANUNCULACEÆ.—The plants belonging to this order are herbs or climbing plants, never shrubs or trees. The flowers are solitary, that is, singly, on a stalk, or in racemes, that is, in bunches of irregular flowrets. The petals are generally five, but sometimes are deformed and very minute, or wanting altogether. The stamens are very numerous, and placed on the receptacle. The fruit is composed of several carpels, distinct or partially united. The seeds are erect or pendulous. The family *Ranunculaceæ* are widely diffused all over the globe, but especially in cool and temperate climates. Within the tropics they are chiefly confined to high mountain districts. They are remarkable for their acrid, poisonous qualities in many of the species of *Ranunculus*; the acrid secretion in the leaves will produce blisters if applied to the skin; whilst the Aconite or Monk's Hood is a deadly poison. A good example of the chief characteristics of the family is afforded by the species figured in our Plate No. 1.

WILD FLOWERS.



BUTTERCUP, OR BULBOUS CROWFOOT.

(Fig. 1.)

RANUNCULUS BULBOSUS.

THE name Buttercup is familiarly applied to nearly all the species of *Ranunculus* with bright yellow flowers. This species is distinguished from the rest by its thickened stem, which at the lower part, under the ground, expands into a sort of bulb, and by the sepals, which, as soon as the flower expands, are reflexed or turned back on the peduncle. The leaves are divided into three stalked segments, more or less cut. The whole plant is about a foot high; it flowers in the early summer, and is abundant in our meadows and waste places. In Scotland it is found southwards; but in the north is seldom to be seen. The genus *Ranunculus* is the type of the natural order *Ranunculacæ*. The species are generally acrid, and not eaten by cattle in their growing state.

TRAVELLERS' JOY—OLD MAN'S BEARD,
OR VIRGIN'S BOWER.

(fig. 2.)

CLEMATIS VITALBA.

CLEMATIS VITALBA belongs to an almost exceptional genus of the *Ranunculus* family. Its stem is climbing, and woody at the base; it is the only British plant which gives us some faint notion of the bush ropes of the tropics. The woody stems sometimes attain a great thickness, and the petioles or leaf-stalks of the young branches act as tendrils, and cause it to spread to a great extent over trees and shrubs in its neighbourhood. The flowers are of a greenish-white colour, in loose bunches. The carpels are very conspicuous when ripe, from their persistent styles, which grow into long feathery awns; hence the name, Old Man's Beard. The petals are absent in this species, the flowers being formed by the sepals. The leaves are pinnate; the leaflets, usually five in number, ovate and slightly pointed in shape. This pretty and slightly sweet-scented plant is one of the greatest ornaments of our country hedges; and we can, doubtless, all remember with pleasure the delight with which in our summer rambles we have torn down long wreaths of its pretty green leaves and pale flowers to adorn a rustic head-dress, or to luxuriate in its fragrance, which it possesses in some degree in common with its relation *C. Flammula*, the deliciously fragrant plant so well known in our gardens.

There are above a hundred species of *Clematis*, most of which are favourite plants in cultivation. Both our own British species, *Clematis Vitalba* and *C. crispa* (an exotic species), have been used as rubefacients in rheumatism, and the dried leaves of *C. Vitalba* form fodder for cattle in some places, the acrid juice they contain when green disappearing after drying. When dried, boys use clematis wood for smoking as they do ratan. A thin slice of the wood is an interesting object under the microscope, from the curious manner in which the parts of the stem are arranged.



WOOD ANEMONE, OR WIND-FLOWER.

(Fig. 3.)

ANEMONE NEMOROSA.

THIS is one of our commonest and prettiest hedge-plants, belonging to the same order, Ranunculaceæ. The whole of the anemones are lowly herbs, usually perennial, as is this species. The flowers are solitary, consisting of six smooth white elliptical sepals. The leaves consist of three ovate or lanceolate leaflets, of a dark bright-green colour. The flowers may be seen in or near woods as soon in the year as April, and our earliest spring walks are often gladdened by its presence. Nothing can be prettier than a bouquet of these simple delicate white flowers in the midst of their natural guardians, the dark finely-cut green

leaves ; they are among the earliest harbingers of the season,

When earth, exulting from her wintry tomb,
Breaks forth with flowers.

To see these delicate flowers in perfection, however, we must choose for our woodland excursion a bright unclouded day, for the Wood Anemone is a natural barometer, and droops at the approach of rain. There are several British species of anemone ; but we have selected the *A. nemorosa* as the most attractive from its very simplicity.

Anemone Pulsatilla, has fine large purple flowers. *A. Pæonia*, the Peacock Anemone, common in the south of Europe, with its bright scarlet or scarlet and white flowers ; and *A. coronaria*, the garden Anemone, with its many varieties of hue, form striking contrasts to the unobtrusive appearance of our little favourite. The whole genus partake of the acrid and poisonous qualities of the family, and are unsafe to take, even as medicines, although they have had a reputation in various complaints.

PHEASANT'S EYE.

(Fig. 4.)

ADONIS AUTUMNALIS.

ADONIS AUTUMNALIS,—Pheasant's Eye,—partakes more of the character of the true *Ranunculus* than

any plant we have yet described ; it is distinguished from that genus by the absence of a little scale at the base of the petals, and from other genera of the same order by the numerous hard, dry, sharp-pointed grains of which its fruit consists. It is an upright annual plant, from eight inches to a foot in height. The leaves are finely cut into numerous narrow linear segments. The sepals are green or slightly coloured ; the petals from five to eight in number, but slightly longer than the calyx, of a bright scarlet colour, with a dark or black spot at the base inside the flower. It is constantly to be found in cornfields in the summer, from May to September in England and Ireland, and sometimes in Scotland.

GLOBE FLOWER.

(Fig. 5.)

TROLLIUS EUROPEUS.

TROLLIUS EUROPEUS, Globe Flower, belongs also to the order Ranunculaceæ, and is so called from the German word *trollen*, round, in reference to the round shape of the flowers. The species are all perennial herbs, with divided leaves and yellow flowers, composed of coloured sepals. It is not a large genus, and our native species, *Trollius Europæus*, is the most generally known. It has from ten to fifteen broad concave sepals, usually turned over in the shape of a ball, and concealing petals, stamens, and carpels.

In Scotland it is called Lucken Gowan, or Cabbage Daisy. In some parts of England, as well as on the continent of Europe, they are gathered on festive occasions for making garlands and decorating the cottages of the peasantry. In common with its natural order, this plant is slightly acrid. It likes a rich, moist soil, but loves a good strong light to flourish under; deriving vigour and colour as the moon derives her light.

COLUMBINE.

(Fig. 6.)

AQUILEGIA VULGARIS.

AQUILEGIA, literally water-gatherer, is another genus of plants of the *Ranunculus* family, so called because the leaves collect water in their hollow. *A. vulgaris*, the Common Columbine of our hedges, is a pretty little plant with a stem from one to two feet high or more. The flowers, which are curious, are of a dull purple or blue colour, and drooping. The root-leaves and those at the lower part of the stem grow in a large tuft, each with a long stalk. The petals have all a long curved horn or spur at their base, which projects below the calyx. The stamens are numerous. The English name, Columbine, is derived from a fanciful likeness to a dove, which is produced if we separate one petal from the flower-cluster; it brings with it two sepals, and the appearance of a dove may be imagined. When wild, the blossom is of a light-blue colour, but

the plant is subject to great changes in cultivation, and readily produces double flowers. We find in "Brown's British Pastorals," that our dove-like plant was, in former times, the insignia of a deserted lover :—

"The Columbine by lonely wand'rer taken,
Is there ascribed to such as are forsaken."

MONK'S-HOOD, OR WOLF'S-BANE

(Fig. 7.)

ACONITUM NAPELLUS.

ACONITUM NAPELLUS is a plant with a firm erect stem one and a half to two feet high, also belonging to the Ranunculaceæ. The leaves are either stalked or very close to the stem, of a dark-green colour and very smooth. The flowers are large, and are easily recognized as having the very large uppermost segment of the calyx overhanging the petals and other parts in the form of a helmet. The two upper petals inside this covering are long and narrow, with spurs; the three lower ones very small. We may all remember having plucked these flowers in our childhood and having thrown back the hood or calyx, have delighted in the fairy chariot and steeds formed by the petals thus set free. No vegetable poisons are more powerful than those produced by this genus of plants; and the common species, *A. Napellus*, yields it in the greatest degree. It is sometimes prescribed medicinally, and according to some writers very beneficially.

From ancient times these plants have been celebrated as virulent poisons. In 1524 and 1526 two criminals at Rome and at Prague, to whom the root was given by way of experiment, very quickly perished. It entered into the deadly draught which the old men of Ceos were condemned to drink when they became infirm, and is also said to have been the principal ingredient in the cup which Medea prepared for Theseus. The most virulent Indian poison, *Bikh* or *Bish*, is supposed to be a preparation of a species of Aconite. Dr. Wallich describes *A. ferox* as used by the native Indians to poison the water in the tanks, in order to impede the progress of a hostile army. It is also used to poison darts, arrows, and spears.

THE WHITE WATER-LILY.

(Fig. 8.)

NYMPHÆA ALBA.

THIS plant belongs to another natural order or family,—Nymphæaceæ, which consists chiefly of aquatic herbs with floating leaves and solitary flowers; found in all temperate and tropical parts of the world. They have usually four sepals and many petals in several rows, contracting gradually into stamens. The fruits are numerous, but are either imbedded into the receptacle, or combined together to form a single ovary with many cells. *Nymphæa alba*, the White Water-Lily, has bright, smooth, heart-shaped leaves,

floating on the surface of the water ; usually six or eight inches in diameter. The flowers are large, white, and floating, with yellow stigmas. It is one of the brightest ornaments of our still lakes and ponds throughout Europe, and is a favourite plant with all lovers of flowers. The flower of the White Water-Lily is an excellent example of the law of morphology in plants. The doctrine that all the parts of a plant are modifications of the leaves, may be aptly illustrated by tracing the gradual changes which take place in the floral envelopes of this plant. Begin with the outermost whorl of sepals, and trace the leaf-like character gradually lessening until they become changed into perfect stamens, with petal-like anthers attached to them. The flower-stems are porous and succulent, but rapidly lose their moisture if removed from the water. The Water-Lily may be transplanted from its native home by placing the thick stems in baskets of earth, and fastening stones to them so as to keep them well under water. These stems have a bitter, astringent taste, but are quite free from any of the poisonous acrid principle of the last family of plants we met with. They have been used in dyeing a dark-brown colour. Goats and swine will eat them, and they have been used medicinally. It is, however, as an object of beauty that the Water-Lily claims our attention ; and nothing can be more lovely than a calm lake on whose bosom may be seen floating numbers of these snowy nymphs. On Loch Lomond acres are covered with them ; and in all the northern English lakes they are more or less abundant. Like the sacred Lotus of the Nile, the flowers rise and

expand as the sun gains strength, and close again in the evening ; sleeping as it were through the hours of darkness until called into life again by the warm rays of light.

Moore poetically describes this natural process,—

“Those virgin lilies all the night,
 Bathing their beauties in the lake,
 That they may rise more fresh and bright,
 When their beloved sun’s awake.”

The stimulus of the sun’s rays seems to have relation to the fertilization of the plant. The pollen, if scattered beneath the water, would be washed away and decomposed, while on the expanded raised flower it is received without injury. This is truly the object for which—

“The water-lily to the light
 Her chalice rears of silver white.”

And as it is with poets in sentiment, so it should be in our every-day life ; each daily duty, if viewed aright, contains in it the elements of poetry, which may be made to surround the most prosaic acts of existence with beauty.

The Yellow Water-Lily, *Nuphar lutea*, is almost as attractive in its golden radiance as is its more modestly-attired sister.

“In golden armour glorious to behold,” it forms a beautiful object on the surface of a lake or river. Other tropical species of *Nuphar* have wonderfully-tinted blossoms, of blue and crimson.

In Demerara grows the *Victoria regia*, whose leaves are as much as six feet and a half across, and

whose flowers are frequently fifteen inches in diameter. All lovers of floral beauty should give themselves the treat of seeing these magnificent flowers in the aquatic greenhouse at Kew, or in the Regent's Park botanical gardens, where, in the season, they blossom in perfection.

PRICKLY POPPY.

(Fig. 9.)

PAPAVER ARGEMONE.

PAPAVER ARGEMONE belongs to the family of Poppies,—*Papaveraceæ*. They are all herbs with much-divided leaves and no stipules. The sepals are two in number, and fall off as the flower expands. The flower consists of four regular-shaped petals. The stamens are distinct and numerous. The fruit is a capsule opening in valves; the seeds albuminous, and containing a fixed oil. *Papaver Argemone*, the Pale Poppy, or Prickly Poppy, is not so common as the Corn Poppy, *P. Rhœas*; but is remarkable for its delicacy. The flowers are of a pale-red colour, with a dark spot at the base; the segments of the leaves are few and narrow, and the fruit or capsule is covered with minute bristles. There are several other species of British poppies: the common Field or Corn Poppy, *P. Rhœas*; the Long-headed Poppy, *P. dubium*; the Rough Poppy, *P. hybridum*; the Opium Poppy, or *P. somniferum*. All the species exude more or less a milky juice, the narcotic properties of which are

considerable only in *P. somniferum*. This species yields the poppy-heads so commonly used in fomentations. Opium can be prepared from the juice of the English plant, which has been used instead of foreign opium ; and although not so powerful in its action, possesses the same properties.

Many are the curious legends and traditions connected with the poppy plant. Theocritus tells us that the silken petals of the poppy prove talismans for Cupid, thus :—

“By a prophetic *poppy-leaf* I found
Your changed affection ; for it gave no sound,
Though in my hand struck hollow as it lay,
But quickly wither'd like your love away.”

In classic lore, the poppy was sacred to Ceres, though our modern notions of agriculture would rather regard it as an intruder into her domains.

Of all the strange and baneful, as well as beneficial effects of the poppy-juice or opium, we can hardly speak here. As a medicine it is most valuable, if carefully administered, and it has often soothed and palliated the sufferings of humanity. As a hurtful and sensual indulgence, its effects are well known ; in Eastern countries, perhaps, better than our own. The fascination of this pernicious habit, and its terrible results, are vividly described in a popular work,—
“Confessions of an English Opium-eater.”

YELLOW HORN-POPPY.

(Fig. 10.)

GLAUCIUM LUTEUM.

THIS plant reminds us of our seaside holiday-time, of sandy beaches and rolling waves ; and as with no objects are mental associations more vivid than with flowers, the sight of a plant but rarely met with, must recall to our minds the circumstances under which we first saw it. Perhaps a first visit to the sea-beach, or to some well-loved spot, and all the delicious refreshment of the sea-breeze, the rest and repose which steal over the most active mind when listening to the tune of the waters, is in a measure brought back again, as we see, even in its dead and dried state, the well-remembered plant of the seashore. Our Yellow Horn-Poppy is the most striking and remarkable of our seashore plants, and cannot fail to arrest the attention of the most casual observer, in a position where so little vegetation flourishes. The foliage is of a pale sea-green colour, which botanists term glaucous : hence its scientific name. It is rough, with short bristles ; the pods or horns are from six to ten or twelve inches long, crowned by the spreading lobes of the stigma. When we see this pretty plant, it is interesting to remember the history of Glaucus, after whom it is named. He was the son of Neptune and of Nais, a sea-nymph, but lived on shore. His nature, however, had some influence on his habits, and he was fond of fishing. One day,

having been very successful in his sport, he laid his scaly prize on a neighbouring marsh, when to his great surprise they began to nibble the green grass, and then

“ Sudden darting o'er the verdant plain,
They spread their fins as in their native main,
Left their new master, and regain'd the sea.”

Amazed at what he saw, Glaucus resolved to test the power of the herbage in his own person ; and no sooner had he bitten it, than his hereditary aquatic propensities seized him, and into the ocean he leaped, when, for his faith and courage, he was received as a denizen among the sea-gods.

In their domain he still shows his royal descent by wearing a golden robe ; and yet, from old affection, high above it he bears his favourite long and curved fishrod, with its point bent, as if a captive fish ever strained it. Glaucus never goes far out to sea, but rather frequents the shores and cliffs ; for Scylla, whom he loved, was turned into a rock, with howling waves around her ; and his faithfulness retains him still close to her side.

The Yellow Horn-Poppy is the “squats” of the Portland Islanders.

DYER'S WOAD.

(Fig. 11.)

ISATIS TINCTORIA.

OUR next specimen belongs to another and very extensive and useful family of plants,—Cruciferæ or

Crossworts. The species are herbs, or rarely undershrubs, with alternate leaves and no stipules; four sepals, four petals of equal size, or two on the outside layer. The stamens are six in number, of which two are generally shorter, or sometimes altogether absent. The fruit is a pod, divided into two cells by a thin partition, from which the valves generally separate when ripe. When long, it is called a silique, and when short, a silicle.

Isatis tinctoria preserves the general characteristics of the family. The stem is from eighteen inches to two or three feet high, branched in the upper part. Its long leaves are smooth and coarsely toothed and stalked, the pods hanging on slender stalks downwards. The flower is of a yellow colour.

The specific name *tinctoria* signifies its use in dyeing or staining. From it was undoubtedly obtained the blue dye or woad with which the Ancient Britons stained their skins.

When the arts of civilized life were not practised, this substance supplied, according to the poet Garth, all the requirements of a fashionable toilette.

“In times of old when British nymphs were known
To love no foreign fashions like their own,
When dress was monstrous, and fig-leaves the mode,
And quality put on no paint but woad.”

At the present day this plant is used by dyers, not on account of its own blue colour, but as a mordant for other colours. Its colouring principle seems to be identical with indigo. It is cultivated in Bedfordshire, Northamptonshire, and Somersetshire.

THE COMMON WHITLOW-GRASS.

(Fig. 12.)

DRABA VERNA.

DRABA VERNA, a cruciferous plant, is one of our earliest spring flowers ; and as soon as the bright days of March and April tempt us out into the fields and lanes, we may look for it on dry walls and banks, with its little stalk of white flowers. It is a small annual plant, and lasts but for a few weeks ; the leaves are all towards the base, ovate or oblong. The seed-pods burst open in the month of May, and scatter the seeds into the ground, where they lie securely till the next season. The tiny flowers droop at night to keep the stamens from the chilly dews of the early spring. The common name, Whitlow-grass, is given on account of the use of the leaves as a poultice in those unpleasant swellings.

It belongs to the genus *Erophila* of De Candolle, a pretty name, expressive of its early appearance.



COMMON WATER-CRESS.

(Fig. 13.)

NASTURTIUM OFFICINALE.

NASTURTIUM OFFICINALE,—also known as *Sisymbrium Nasturtium*, the Common Water-cress,—should

13



Water Cross

14



Rock Rose

15



Sweet Violet

16




Succow

17



Clove Pink

18



Red Campion

19



Fleck

20



Tree Mallow

21




Music Mallow

22



... Mallow

23



St. John's Wort

24



... Papaveris

be familiar to every one, as belonging to the family Cruciferæ. It is very likely to be confounded with a poisonous plant with which it grows, the Fool's Cress, as it is called (*Sium nodiflorum*). From this it may always be distinguished, and, in fact, from all other Umbelliferæ, by the petioles of the leaves not forming a sheath round the stem. In addition to the characters of the genus, the Water-cress is known principally by the form of its leaves. The leaf is composed of from five to seven leaflets, which are arranged opposite each other on a common leaf-stalk with a terminal leaflet. The leaflets are somewhat heart-shaped and slightly waved or toothed; they are succulent and their surface is smooth. The terminal leaflet is always the largest. The petiole or leaf-stalk does not in any manner embrace the stem. The flowers are white, and the pods, when ripe, are about an inch long. It is a native of rivulets throughout the world, and is very plentiful in our own country. The ancient reputation of this plant as an article of food, valuable both for its pleasant pungent taste and its antiscorbutic properties, is well founded. Recent writers on the subject of diet have shown that in partaking of fresh uncooked vegetable food in the shape of salads or fruit, we are obtaining those salts of potash and other constituents so necessary to health, which in the process of cooking are dissolved away.* Water-cresses are said to contain iodine.

No better vehicle for the introduction of these

* See Dr. Lankester's Lectures on Food. London: R. Hardwicke.

important substances can there be than fresh bright water-cresses ; and our old friend Gerarde's notion of their value presages all the modern discoveries as to their virtue. He says that the eating of water-cresses restores their wonted bloom to the cheeks of sickly young ladies. He might have added that a walk to the running stream where they grow would enhance the effects of the remedy.' So large is the consumption of water-cresses in London that they are cultivated by market-gardeners to a great extent by means of artificial water-supplies ; but none are so delicious as those from natural streams. Our popular street-cry has been rhymed by Swift thus :—

“Fine spring water-grass, fit for lad or lass.”

The name *cress* has, according to writers, many origins. One says it signifies water *cross*, from its cruciate flowers. Chaucer employs the Saxon word *kers* (*cress*) to signify anything worthless :—

“Of paramours ne raught he not a kers.”

From which, perhaps, is derived the phrase of not caring a curse for a thing. *Nasturtium* is a name given to all these biting plants, each being a *nasus tortus*, or nose-twitcher. Pliny records that they put the nose into convulsions. Long ago the refreshing nature of these plants as food was recognized, and there is a Greek proverb, “Eat cress to learn more wit.” Knowing, as we now do, the influence of the physical over the mental, we can rationally understand the proverb.

There are several other species of *Nasturtium*

native in England ;—the Creeping Cress, *N. sylvestre* or *N. palustre* ; the Water-rocket, *N. terrestre* ; and the Water-radish, *N. amphibium*.

THE ROCK ROSE, OR ROCK CIST.

(Fig. 14.)

HELIANTHEMUM VULGARE.

OUR next plant belongs to a new family,—Cistaceæ. The characters of the order are that the species are shrubs or herbs with opposite, or in a few exotic species, alternate leaves, with or without stipules, generally smelling fragrantly. The petals are usually five in number, broadly spreading, the sepals three, nearly equal, overlapping each other in the bud, with or without two smaller outer ones.

The species figured *H. vulgare*, is a low under-shrub, with a woody stem ; the leaves have stipules ; the flowers are of a bright-yellow colour, broadly spreading, and blooming from May to September. It is found in dry meadows and pastures throughout Europe and western Asia, and is not uncommon in Great Britain. The Rock Rose, or Cistus of our gardens, is a variety of this species.

SWEET VIOLET.

(Fig. 15.)

VIOLA ODORATA.

THE SWEET VIOLET is a favourite with everybody, and scarcely requires description to be recognized. It is, however, interesting to know that it belongs to the family *Violaceæ*, and to the only European genus of that family. It has five petals of unequal shape and size, the lower one being drawn out into a kind of spur. There are five sepals, and the stamens are connected together; two of them with curious ear-like appendages. The flowers are of a purplish colour, nodding. On the stem we have an example of what are called bracts. The leaves grow at the base of the plant, with rather long stalks, and are broadly heart-shaped. There are several British species of this genus, but our sweet-scented violet, and the pretty blue dog-violet, which is inodorous but very attractive, are those most worthy of notice. Who does not welcome the first violets in the early spring, "gleaming like amethysts in the dewy moss;" and there is no land where these pretty flowers grow in which their praises have not been sung. We must all have felt the power of perfumes in recalling to the memory images and scenes of past years, before these lines were written,—

"The smell of violets hidden in the grass
 Poureth back into my empty soul and frame
 The times when I remember to have been
 Joyful and free from blame." TENNYSON.

Not only is the violet celebrated for its beauty, but for its uses and for its mystic powers. Violet roots and violet flowers have been used as remedies in all sorts of diseases. The Athenians were noted for their love of these flowers, and they were reputed to "moderate anger," to procure sleep, and to comfort and strengthen the heart. At the present time the root is used as an emetic. Pliny prescribes a liniment of violet roots and vinegar for gout and disorders of the spleen. The violet is certainly a classical plant. It was a favourite with the old Greeks. Homer and Virgil both mention it frequently, and Shakspeare alludes to a very old superstition, when he says,—

"Lay her i' the earth,
And from her fair and unpolluted flesh
May violets spring."

SUNDEW.

(Fig. 16.)

DROSERA ROTUNDIFOLIA.

A VERY pretty and curious little plant is the Sundew, or *Drosera rotundifolia*; and it is found where we should least look for beauty: in bogs and morasses, in the damp corners of heaths and wildernesses, we descry the ruby points of the leaves of this lovely little plant sparkling amid emerald-green moss-tufts. It is the type of the family Droseraceæ. A small flower-stalk, from two to six inches in height, bears on

the top the few little white flowers which expand in the sunshine. The leaves grow very low down, close to the ground, and are of a round shape, and thickly covered with the minute red hairs, each of which secretes a drop of fluid, which sparkles in the sunshine like diamonds. These drops of fluid are of a somewhat glutinous nature, and are apt to entrap unwary insects that happen to alight on the leaves. It is somewhat difficult to find the Sundew fully expanded; the best way of seeing it in all its beauty, is to remove a tolerably large portion of the plant with a trowel, and then, placing it in a saucer, surrounded with the damp green moss in which it grows, cover it over with a hand-glass, supply it freely with water, and you have as pretty a little Ward's case as can be imagined, and an interesting subject for botanical study likewise. It is to be found, by careful searching, on Hampstead Heath, on Wimbledon Common, and in most boggy, heathy districts. It is an acrid and caustic plant, and has been supposed to cause the rot in sheep. It had once a reputation as a cosmetic, and is said to burn away corns and warts. It also curdles milk, and has been frequently used in the dairy for that purpose. This is the plant of which Burton, in his "Anatomie of Melancholy," says "that Bernardus Ponottus prefers his herba solis before all the rest of herbs in this disease (melancholy), and will admit of no herb upon the earth to be compared with it." In Italy it is used for making the liquor called Rossoli.

There are other species of *Drosera* in the British Flora, which possess dyeing properties, as may be

seen when they are dried in the herbarium, by the red colour they communicate to the paper.

THE CARNATION, OR CLOVE PINK.

(Fig. 17.)

DIANTHUS CARYOPHYLLUS.

THIS plant is a native of the South of France, but is found wild on old walls in Kent and Norfolk. It belongs to the family Caryophyllaceæ, and is one of a beautiful genus of plants, all of which are sweet-scented and showy. Mr. Babington describes six species of *Dianthus* as natives of Great Britain. The clove pink has solitary flowers; the scales of the calyx are much shorter than the tube of the corolla; the leaves have smooth margins; the petals are toothed, ovate, and smooth; the stem branching, long, and procumbent; the seeds nearly flat, and the flower-stem from twelve to eighteen inches high. The flowers of the genus are of all colours, excepting blue; in the British species they are of a pale pink, and in all cases fragrant. The flowers of the Clove Pink are used to give colour and fragrance to a syrup used in medicine. A small variety of this species is known by the name of Picotees.

THE RED CAMPION.

(Fig. 18.)

LYCHNIS DIURNA.

THE RED CAMPION, *Lychnis diurna*, belongs to the family Caryophyllaceæ. It has oblong leaves, usually pointed, tapering to the base, the lower ones stalked. The flowers are few, in loose branches, of a red colour, opening in the morning; the calyx forming a sort of capsule composed of two divisions, or teeth, which curve backwards. The capsule thus formed becomes globular as it ripens. This pretty plant is found in moist shady woods and hedge-banks all over Britain. It flowers all through the summer, beginning early in the spring. The other British species are known by the names of Scarlet Lychnis, Ragged Robin, and White Champion. The Corn Cockle is also a pretty plant belonging to this family, which is found blossoming in cornfields from June to September.

FLAX.

(Fig. 19.)

LINUM USITATISSIMUM.

ONE of the most valuable of our native plants is the Flax Plant,—*Linum usitatissimum*; and in the

delicate little blue flowers and fragile-looking stem the uninitiated would little imagine the living of thousands and the chief manufacture of a great country to lie concealed. It is a tall, erect annual plant, with alternate narrow-pointed leaves, and flowers of a rich blue colour, arrayed in a loose terminal bunch. The fibre of the plant is woven into linen, and the seeds are valuable on account of the mucilaginous nature of their external coating, and for the oil they contain, which is prized for burning and as a drying medium in the arts; the oilcake made from the seeds is an extensive article of food for cattle. Flax is chiefly cultivated in Ireland, and from a very early time the importance of its culture was recognized in the economy of this country. In 1750, Sir William Temple wrote a treatise on the subject.

Microscopic examination has clearly proved that the mummy-cloth of Egypt is made of linen fibres, and not, as was once supposed, of cotton. Herodotus and Plutarch tell us that it was not permitted to any Egyptian priest to enter a temple unless he wore a linen garment. By the Greeks, linen was used in the time of Herodotus; possibly not produced by the same species as the one now grown for the purpose, but undoubtedly the fibre of a species of *Linum*. We have four British species of this genus. None are of any importance but the one we have selected. The Welsh species, or the fairy flax, is an elegant little mountain species, which is remarkable for its beauty and grace, and has a medicinal reputation in its native mountain districts. The exquisite deli-

cacy of the flax plant is aptly pictured by Coleridge :—

“The unripe flax,
When through its half-transparent stalk at eve
The level sunshine glimmers with green light.”

THE TREE MALLOW, OR SEA LAVATERA.

(Fig. 20.)

LAVATERA ARBOREA.

THE genus *Lavatera* was named in honour of Lavater (a physician in Zurich, not the celebrated physiognomist of that name), and belongs to the Mallow family,—*Malvaceæ*. The flowers in all the genera of this family are twisted in the bud, the calyx composed of five divisions, with three or more bracts at the base, forming an outer calyx. The species we are now considering has a woody stem, something like that of a cabbage, with thick, hard, flowering branches. The leaves are on long stalks, with seven, five, or three lobes, and soft as velvet. The flowers are mostly in pairs, of a pale purple-red colour, with dark blotches at the base of the petals. It blossoms from July to October, and is found on seaside rocks throughout Europe; in Britain, chiefly on the south and west coasts of England and Ireland, especially in the Isle of Wight, and on the Bass Rock in the Frith of Forth.

THE MUSK MALLOW.

(Fig. 21.)

MALVA MOSCHATA.

THIS plant belongs to the family Malvaceæ. The genus Mallow is a very favourite and well-known group, and most of us may recall, when children, the delight with which we have sought for the plants, in order to secure the much-prized "*cheeses*," or fruits of the mallow. The Musk Mallow has a tough woody root, and an upright partially-branched stem. The root-leaves have long leaf-stalks, with rounded limbs cut into three or five lobes. The stem-leaves are much more deeply cut and lobed than the root-leaves. The whole herbage is more or less hairy. The flowers are large, rose-coloured, or rarely white, crowded together at the top of the stem and branches. The calyx consists of two whorls, the outermost composed of three acute sepals, the innermost five-toothed. The corolla has five wedge-shaped petals jagged at the end. Were this pretty plant less common than it is, it would, perhaps, meet with more admirers; but the extraordinary perverseness of the human mind leads us to admire most that which is least attainable, and to hold an everyday object in slight estimation. The musk-like scent of the particular species of mallow we have selected distinguishes it from the rest, although it be but slight. The Musk Mallow, like the anemone, closes its petals at night. In floral

language it is regarded as the emblem of a sweet, mild disposition, and we read that it was customary with the ancients to plant musk mallows around the graves of their departed friends. The common mallow, *M. sylvestris*, yields, when boiled, a plentiful tasteless mucilage, which is used in some cases as a medicine.

MARSH MALLOW.

(Fig. 22.)

ALTHÆA OFFICINALIS.

THIS species is also a Malvaceous plant, but belongs to a different genus. Marsh Mallow,—*Althæa officinalis*, is a common European plant, and is often found in marshes, especially near the sea, in great abundance. It is a perennial, with a carrot-shaped, white fleshy root, as thick as the thumb, and a foot or more long. The stems are two or three feet high, covered all over with a soft down, which is also found on the leaves, and gives them a hoary appearance. The leaves are a little heart-shaped and three-lobed, The flowers are not large, of a pale rose-colour, and appear in very short clusters from the bosom of the leaves; the calyx is five-toothed, and surrounded with bracts. The whole plant, and particularly the root, abounds in mucilage; the demulcent lozenges sold in the shops under the name of *Pâte de Guimauve*, are made of marsh mallow, and a few years ago there was an ointment commonly used, which may still be in repute for aught we know, called Marsh Mallow Oint-

ment. The hollyhock of our gardens is a species of *Althæa* from the Mediterranean. It is found wild in China.

THE LARGE-FLOWERED HYPERICUM,
OR ST. JOHN'S WORT.

(Fig. 23.)

HYPERICUM CALYGINUM.

THIS is a showy plant, belonging to the family *Hypericaceæ*. It was brought to England by Sir G. Wheeler in 1676, from its native woods near Constantinople. It has been long cultivated in our gardens, and is found wild in bushy places in England and in Ireland. This species of St. John's Wort has a creeping woody rootstock, the stems scarcely a foot high, simple, and branching at the base only, with large oblong leaves, green and smooth, having very small clear spots on their surface. The flowers are of a bright yellow colour, three or four inches in diameter, with one or two on the top of each stem; in our gardens it is more luxuriant, and produces five or six. There are thirteen British species of this genus, and one hundred and seventy-two are enumerated by De Candolle as growing in various parts of the world. They are all known as St. John's Worts, and most of them are worthy of cultivation. The hardy herbaceous kinds will thrive in any common garden-soil, and are easily propagated by dividing the roots or by seeds. Those

which require the greenhouse will thrive best in a mixture of loam and peat, and strike root readily under a hand-glass.

The *Hypericum perforatum*, the perforated St. John's Wort, was one of the flowers gathered by our forefathers to be thrown into the bonfires which were kindled in London on the eve of St. John. In many parts of France and Germany the peasants still gather its golden blossoms, and hang them with much ceremony in their windows and doorways, as a charm against evil spirits, storms, thunder, and all calamities. This custom probably arose from the misinterpretation of some medical writers, who, believing in its virtue as a remedy in maniacal disorders, called it *Fuga Dæmonum*. It was at one time worn about the person in Scotland as a protection against witchcraft and enchantment.

GRASS OF PARNASSUS.


(Fig. 24.)

PARNASSIA PALUSTRIS.

HERE we are among the gods; but surely this pretty plant has mistaken its habitation, for we find it in marshy, boggy places, and scarcely ever on elevated land. The Grass of Parnassus, or *Parnassia palustris*, belongs to the same family of plants as our little Sundew,—*Droseraceæ*; and it is often found in the same localities. Why it is called a grass we cannot



25



Red Crane's Bill

26



Wood Sorrel

27



Rest Harrow

28



Strawberry clover

29



Yellow Vetchling

30




Creeping Cinquefoil

31



Willow Herb

32



Evening Primrose

33



Purple Loosestrife

34



Wild Bryony

35



House Leek

36



Marsh Marigold

imagine ; for with its delicately-coloured flowers, beautifully netted, it does not resemble the family of grasses at all. The leaves around the roots are on long stalks, while here and there a leaf clasps the flower-stem. Each flower-stalk produces but one flower, which has curious fringed scales, which lie around the centre, and are the nectaries ; a little waxlike gland exists at the tip of each hair. In former times, this flower was called the Noble Liverwort, and was doubtless considered a remedy in diseases of the liver. It is the only British species of the genus. Sir William Hooker tells us that this elegant plant, if plunged into water in a garden-pot, surrounded by a ball of its native earth, will continue to blossom for many weeks.

RED CRANE'S-BILL, OR THE BLOOD
GERANIUM.

(Fig. 25.)

GERANIUM SANGUINEUM.

THE BLOOD GERANIUM grows in great profusion on our western limestone coasts, and is one of the most attractive of the species. The family to which it belongs is Geraniaceæ ; of which the characters are well preserved in the genus Geranium.

The plants known by this name must not be confounded with the so-called geraniums of our gardens, which, though of the order Geraniaceæ, belong to the

subdivision *Pelargonium*. The five petals of our true geranium are of the same size, while the inferior petals of the pelargoniums are smaller than the other two, and of a different character.

Geranium sanguineum is known by its numerous stems, about a foot long, leaning downwards, or sometimes upright, with spreading hairs. The peduncles are mostly single-flowered; the carpels smooth, crowned with a few bristles; the leaves nearly round, seven-lobed. The flowers are large, of a dark-purple colour; the sepals hairy, with a fine point. The whole plant somewhat resembles the mallow in appearance, and there is an Eastern notion that geraniums were at first simply mallows, until Mahommed, delighted with the fine texture of a shirt made for him of mallow fibres, turned that plant into the more beautiful geranium; or some assert that his shirt being spread to dry on a mallow plant, the transformation was discovered on taking it up. There are above a dozen British geraniums; the most common one, probably, being *G. Robertianum*, the Herb Robert, which has a medical reputation. The root of *G. maculatum*, or Alum-root, is a really powerful astringent, containing more tannin than catechu or kino. In North America it is employed as a remedy in sore throat among children. The beautiful red colour assumed by the leaves in dyeing may have been suggestive of their efficacy in arresting hæmorrhages; the fancy is scarcely more absurd than our modern doctrine "that like cures like."

THE WOOD SORREL.

(Fig. 26.)

OXALIS ACETOSELLA.

THE WOOD SORREL, *Oxalis acetosella*, belongs to the same family as the Geranium,—*Geraniaceæ*. It is easily recognized by its three delicately green leaflets, with longish stalks, marked with a darkish crescent in the centre, veined, and its lovely white flowers, which at first sight resemble the wood anemone. There are few walks or shady woods where, in the early spring, the bright half-folded green leaves of this pretty little plant may not be found. The tiny white flowers, with their delicate purple veins, are called by the Welsh “fairy bells,” and are believed to ring the merry peals which call the elves to “Moonlight dance and revelry.” The whole plant abounds in an acid cooling juice, which contains oxalic acid. An infusion of the leaves is frequently administered as a cooling drink in Russia, and the salt prepared from it is used under the name of “salt of sorrel,” to remove the stains of ink and iron-mould from linen. Among the Druids, its triple leaflets were regarded as a mysterious symbol of a trinity, the full meaning of which was involved in darkness. So, too, St. Patrick chose this leaf as his symbol to illustrate the doctrine he sought to teach, and converted many by the apt use of an illustration derived from a plant already sacred in the eyes of his hearers. The original *shamrock* was

undoubtedly the *Oxalis*; though the name became applied to all sorts of trefoiled plants. In all ages, the trefoiled leaf has been regarded with veneration, especially when, departing from its usual form, it is found with four leaflets. The old song, "I'll seek a four-leaved Shamrock," tells of the fairy spells to be woven with its enchantment.

The wood sorrel approaches nearest of all our native plants to a sensitive plant, not only closing its petals and folding its bright green leaves at sunset, and with every change of atmosphere, but even if the stem be rudely or repeatedly struck. In a flower-saucer, or pan covered with a glass shade, this pretty plant forms a lovely object; and we have frequently seen it blossoming year after year in a Ward's case or under a hand-glass, covering the space given to it with its delicate white blossoms. An old Welsh proverb says—

"Three things let no one trust such as shall dislike them,—
The scent of trefoils, the taste of milk, the song of birds."

THE REST-HARROW.

(Fig. 27.)

ONONIS SPINOSA.

THIS plant belongs to the Pea-flower family, or Leguminosæ. It is a low spreading little shrub, with hard, tough, trailing roots, which often retard the progress of the plough, while its numerous and thorny

branches are so great an impediment to the action of the harrow in turning up the ground, as to have obtained for it the common English name of Rest-Harrow. Frequently the whole plant is clothed with sharp hairs or prickles, but sometimes they are quite soft, and the plant smooth, according to the soil in which it grows. The leaflets are oblong or obovate. The flowers are solitary, on short branches of a pink colour, sometimes streaked with a darker shade. It is often very abundant on the sandy cliffs near the sea-shore. This little plant has its uses; for although in its thorny state no animal but the donkey will eat it, yet on better soils it mingles with the pasture, and is relished by cows, sheep, and goats. The roots are succulent and sweet, and have been considered good as a pickle. In the time of Queen Elizabeth they were often brought to table in this form.

THE STRAWBERRY CLOVER.

(Fig. 28.)

TRIFOLIUM FRAGIFERUM.

EVERY one must have noticed, and perhaps gathered, this plant, wondering whether it were fruit or flower, scarcely recognizing in the tiny red heads, the closely-packed flowers surrounded by an involucre of bracts. It belongs to the family Leguminosæ, and is found chiefly in damp meadows and in moist places near the sea. In the neighbourhood of London it is not

uncommon. There are many British species of clover. Mr. Babington describes twenty-one. The white or Dutch clover is very largely cultivated, and is now often used as the representative of the true shamrock, or three-leaved plant of Ireland. The original mystic plant was doubtless the *Oxalis*, of which we have before spoken. Leaves of this character have from a very remote period been regarded with superstitious veneration.

“The holy trefoil’s charm”

was considered potent against all manner of evil.

THE YELLOW VETCHLING, OR YELLOW PEA.

(Fig. 29.)

LATHYRUS APHACA.

THIS plant is a common European field plant, belonging to the natural order Leguminosæ. It is a little smooth pale-green annual, branching from the root into several weak stems, either lying on the ground or climbing by means of numerous alternate simple tendrils, each of which springs from between a pair of large stipules of a broad arrow-shape. There are no true leaves or leaflets, except now and then near the root. The flowers are solitary on simple stalks, small, drooping, and lemon-coloured. The bracts are in pairs, oval-shaped; the teeth of the calyx long

and lanceolate, ribbed. The pod is about an inch in length, smooth, and containing about six round seeds, which are somewhat narcotic, and produce excessive headache if eaten abundantly when ripe. In their young and green state they may be eaten like green peas without inconvenience. *L. sylvestris*, the everlasting pea, is a well-known and pretty species of the same genus. It is cultivated in our gardens, and is often found wild in England. The genus *Lathyrus* is nearly allied to *Vicia*, the true vetches, of which there are many very pretty British species; *Vicia cracca*, the tufted or purple vetch, is one of the most attractive. We can imagine that it must have been this pretty flower which suggested to Sir Walter Scott these lines:—

“And where profuse the wood-vetch clings
 Round ash and elm in verdant rings,
 Its pale and azure pencill'd flower
 Should canopy Titania's bower.”

CREEPING CINQUEFOIL.

(Fig. 30.)

POTENTILLA REPTANS

THIS is a common British plant, belonging to the family Rosaceæ, and is found on heaths, moors, and open pastures throughout Europe. The stems are prostrate, and creep along the ground, rooting at the joints for a considerable distance. The stipules are ovate, and mostly entire; the calyx hairy. The

leaves are stalked with five obovate or oblong coarsely-toothed leaflets. The petals are large, of a bright yellow colour, mostly five in number, sometimes four. This pretty plant is found on the borders of meadows, edges of woods and hedges throughout Europe and Asia. It greatly resembles the common *Tormentilla*, and is often mistaken for it. The latter is an astringent plant, and has been used medicinally; also for tanning in the Western Islands of Scotland and the Orkneys.



WILLOW-HERB, OR CODLINS AND CREAM.

(Fig. 31.)

EPILOBIUM ANGUSTIFOLIUM.

APPLE-PIE PLANT, as it is called, is a very handsome plant, belonging to the family Onagrarix. It has a creeping root, an erect nearly simple stem, slightly hoary, but not hairy, like another species, *E. nilobium hirsutum*. The leaves are shortly stalked, lanceolate, entire, or with very minute distinct teeth. The flowers are large, purplish red, in long bunches; the petals are slightly unequal, entire, and spreading from the base; the stamens and styles inclined downwards. The stigmas are deeply four-lobed. The pod is from one to two inches long, more or less hoary. It is found in mountains, woods, and meadows, in Europe and Siberia. In Great Britain it is found in moist

places in the north of England, and in the south of Scotland. It has crimson, inodorous flowers, with blue pollen. It is a showy plant, and is often transplanted into gardens, where, however, it must be carefully watched, or its creeping roots will encroach on other plants. The down of the seeds furnishes that soft downy substance which, either alone or mixed with cotton, is often woven into stockings, gloves, and such things. The leaves are used in the adulteration of tea, or as a substitute for it. Its young root-stalks and suckers are boiled and eaten; and the Kamtschatskans make a beer from an infusion of the plant. The pith is dried and boiled, and on being fermented is converted into vinegar.

The name Willow-herb is given probably from some slight resemblance in the outline of the leaves to those of a species of willow; and perhaps, too, the situations in which the greater part of the species grow being near the water, or in it, may account for it.

Gerarde says the willow-herbs stop bleeding, heal wounds, and drive away snakes, gnats, and flies.

THE EVENING PRIMROSE, OR COMMON ŒNOTHERA.

(Fig. 32.)

ŒNOTHERA BIENNIS.

ŒNOTHERA BIENNIS belongs to the same family as the willow-herb,—*Onagrarizæ*. It is a biennial, with

ovate, lanceolate, flat, toothed leaves, a rough hairy stem ; the petals longer than the stamens, and about half as long as the tube of the calyx. The flowers are large, numerous, and of a bright yellow colour, emitting a slight fragrance. They close in the bright daylight, and open themselves as the sun goes down : hence their name. It is found abundantly on the Lancashire coast, and covers several acres of ground near Woodbridge in Suffolk. The Suffolk poet Bernard Barton has immortalized this, his native flower, in some beautiful lines, which we cannot refrain from quoting here :—

Fair flower that shunn'st the glare of day,
 Yet lov'st to open, meekly bold,
 To evening's hues of sober grey,
 Thy cup of paly gold.

I love to watch, at silent eve,
 Thy scatter'd blossoms' lonely light,
 And have my inmost heart receive
 The influence of that sight.

I love at such an hour to mark
 Their beauty greet the night-breeze chill,
 And shine 'mid shadows gathering dark
 The garden's glory still.

For such 'tis sweet to think awhile,
 When cares and griefs the breast invade ;
 In friendship's animating smile,
 In sorrow's dark'ning shade.

Thus it bursts forth like thy pale cup,
 Glistening amid its dewy tears,
 And bears the sinking spirit up
 Amid its chilling fears.

But still more animating far,
 If meek religion's eye may trace,
 E'en in thy glimmering earth-born star
 The holier hopes of grace.

The hope,—that as thy beauteous bloom
 Expands to glad the close of day,
 So through the shadows of the tomb
 May break forth mercy's ray.

The pretty plant which suggested to the mind of our Quaker poet these beautiful thoughts is the only native species. It is sometimes called the Tree Primrose. The roots are eatable, and were formerly taken after dinner to flavour wine, as olives now are; therefore, the genuine name was changed from *Onagra*, the Ass Food, to *Ænothera*, the Wine-trap. We are not sure whether the change was necessary for such as need an incentive to imprudent potations.

PURPLE LOOSE-STRIFE.

(Fig. 33.)

LYTHRUM SALICARIA.

THIS is a showy plant, belonging to the family Lythraceæ. It has a perennial root-stock, with short annual erect stems, two or three feet high, slightly branched, glabrous, or softly downy. The leaves are of a dark-green colour, opposite, or sometimes in threes, sessile, clasping the stem at the base, lanceolate and entire, from two to three inches long. The flowers are of a reddish-purple or pink colour, in whorled, leafy spikes. It is the *Lysimachia* of Pliny,

a name which is applied to the Yellow Loose-Strife of our hedges. Our word Loose-Strife is a simple translation of this Latin word. A strong decoction of this plant acts as an astringent, as it contains tannic acid.

WILD BRYONY, OR RED BRYONY.

(Fig. 34.)

BRYONIA DIOICA.

A PLANT belonging to the Cucumber family,—Cucurbitaceæ, must not be confounded with the Black Bryony, *Tamus communis*, which is a very different plant, and about which we shall hear presently. It has a thick, tuberous, perennial root-stock, sometimes branched, the annual stems climbing to a great length; it is rough with minute hairs, containing an acrid juice, and emitting a sickly smell when drying. The tendrils are simple or branched, and spirally twisted. The leaves are divided into five or seven coarsely-toothed lobes, of which the middle is the largest. The barren and fruitful flowers are on the same plant, but on different stalks. The barren flowers are placed in small bunches, several together, and are of a pale-yellow colour; the fruitful flowers are much smaller, generally two together, nearly round. The berries are red or orange, about four lines in diameter, containing several flat seeds.

It is a very common plant in hedges and thickets in England, but it is not found in Scotland and Ireland.

HOUSE LEEK.

(Fig. 35.)

SEMPERVIVUM TECTORUM.

HOUSE LEEK, *Sempervivum tectorum*; so named from its tenacity of life; from *semper* and *vivo*, always living. The little plant known by this name belongs to the family Crassulaceæ. There is but one British species. The foreign varieties are some of them very elegant, and are grown in gardens and greenhouses. Our own species was originally a native of the Alpine and sub-Alpine regions of Central Europe, but it has now found its way to the tops of old walls, and the thatched and tiled roofs of the houses of all the countries of Europe. It is frequently called Jupiter's Eye, Bullock's Eye, or Jupiter's Beard. It has very thick and fleshy leaves, the lower ones, one to one inch and a half long, ending in a small point, and bordered by a line of short stiff hairs. The flowers are pink, arranged along the spreading branches of the cyme, without stalks. The petals are twelve in number, smooth within, fringed with delicate hairs at the edges and on the outside.

This plant is very closely associated with the Stonecrops or Sedums, and shares with them virtues both supernatural and physical. It is considered lucky by the Welsh peasantry to have their roofs covered with these plants; they are believed to protect the house from the ravages of the elements, and to insure the

prosperity of the inmates. Pliny mentions the stone-crop as infallible for procuring sleep ; but to produce this effect the plant must be wrapped in black cloth, and carefully introduced under the pillow of the patient without his knowledge. The juice of the House Leek, when mixed with cream or applied by itself, is said to give relief in burns and other external inflammations. Its vitality is such, that it survives the longest droughts, and the rapidity with which it reproduces itself, even after the roughest treatment, is surprising. Its power of continuing to live under the most adverse circumstance is owing to the facility with which it abstracts carbonic acid gas from the atmosphere. This substance is the food of all plants, and it is thus that they take out of the air and appropriate to their own nourishment what would be injurious to man.

THE MARSH SAXIFRAGE.

(Fig. 36.)

SAXIFRAGA HIRCULUS.

THE SAXIFRAGES are called also Stone-breaks, from the wonderful manner in which the fibres of their roots penetrate the strongest rocks and most unpromising soils. The genus *Saxifraga* is the type of the family Saxifragaceæ. Our example is the Yellow Marsh Saxifrage. The stem is solitary, from four to eight inches long, sometimes covered with rusty hairs.

The stem has a purplish hue, and bears but one rather large yellow flower with red dots. The leaves are alternate, narrow, oblong or linear, and entire. This is one of the rare plants which it is so interesting to find. It inhabits wet, turfy moors, at high elevations, and has not yet been supposed to extend further north than Berwick. It is worth cultivation, and will grow well in peaty bog earth. The saxifrages of Great Britain include the London Pride, that pretty well-known flower of the gardens, which is the *S. umbrosa*.

The *S. oppositifolia*, or Purple Saxifrage, is a favourite spring plant, which, though found growing in wild luxuriance on the Welsh and Highland mountains, is eagerly sought for and sold in Covent Garden Market as an early spring flower. There are many other British species. Mr. Bentham describes thirteen. Medicinal properties have been attributed to some of them, and the white flowers of the common saxifrage were supposed to indicate that it was "governed by the moon."

THE WATER PARSNIP.

(Fig. 37.)

SIUM ANGSTIFOLIUM.

THE narrow-leaved Water Parsnip is a plant belonging to the family Umbelliferæ. It has an erect stem, branched and leafy. The leaves pinnate, the leaflets unequally lobed and serrated; the umbels or

branch of flowerets, opposite the leaves. It is found in wet ditches and shallow streams, and being of a poisonous nature, ought to be carefully distinguished from the water-cress, with which it often grows. Another species, *S. nodiflorum*, is, however, more frequently mistaken, and is therefore called Fool's Water-cress.

THE GOLDEN SAXIFRAGE.

(Fig. 38.)

CHRYSOPLENIUM ALTERNIFOLIUM.

THIS plant belongs to the order Saxifragaceæ, but is not certainly a true Saxifrage. The habit of thus giving common English names to plants of different genera leads to much confusion. It has alternate leaves, the lower ones kidney-shaped, hairy, and seated on long stalks. The flowers are small, in little branches surrounded by leaves, and of a deep-yellow colour. It is not a common plant in Great Britain, although pretty generally distributed. It flowers in the spring by the sides of rivulets, and in moist shady places.

MARSH PENNYWORT.

(Fig. 39.)

HYDROCOTYLE VULGARIS.

AN UMBELLIFEROUS plant, having a slender perennial stem, creeping along the wet mud, or even

37



Water Parsnip

38



Golden Saxifrage

39



Marsh Pennywort

40



Sea Holly

41



Samolus

42



Wild Parsnip

43



Mistletoe

44



Elder

45



Honeysuckle

46



Yellow Bedstraw

47



Fuller's Teasel

48



Chocory

floating in water ; rooting at every joint, and sending out from the same point small tufts of leaves and flowers. The leaves are orbicular, one to one inch and a half in diameter, smooth, and attached by the centre to rather a long stalk. The flowers are very small, white, and on short stalks. The shape and size of these leaves somewhat resemble a piece of money ; hence the name pennywort. It is also known as Pennygrass, White-rot, Fluke-wort, and Sheep's-bane. These latter names it has obtained on account of its being supposed to produce the rot and other diseases in animals that feed upon it. This is, however, an error, as it does not produce disease, but occurs in damp, moist situations, where animals that feed are likely to be attacked with rot and other diseases.

SEA HOLLY.

(Fig. 40.)

ERYNGIUM MARITIMUM.

I AM glad to write again of a sea-shore plant, for I associate with them quiet bright holiday hours ; the never-weary sea, rippling on the clean smooth beach ; happy children with spades and willing hands, making fairy gardens in the sand, filled with the wild flowers which grow around and up the sides of the cliffs, whither they climb, regardless of downfalls, to secure their treasures ; but woe to the little hands that recklessly seize in their grasp the treacherous Sea Holly, with its stiff, sharp-pointed, prickly leaves.

They are covered with a bluish or sea-green bloom, being what is called in botany, glaucous. This character is very common in plants growing near the sea, but not confined to them. One would not easily recognize the Eryngium as an Umbelliferous plant ; but if it is carefully examined, it will be found to preserve all the characteristics of the order. The upper leaves embrace the stem, which is about a foot high, and are lobed and palmate in shape. The flowers are in heads rather than umbels, of a pale-blue colour. It is very abundant on the eastern shores of England, and is found in Scotland and Ireland. The plant is often called Sea Eryngo, Sea Hulver, and Sea Holme. According to Linnæus, the flower-shoots are very good when boiled and eaten like asparagus. The leaves are sweetish, with a warm aromatic flavour. The root also is sweet to the taste, and has a warm, aromatic smell. It has been used in medicine, and was recommended by Boerhaave, the great Danish physician. It is candied and sold in the shops in London as a sweetmeat. At Colchester, in Essex, there exists an establishment where this preparation was first made, more than two centuries ago, by Robert Buxton, an apothecary. There is also another British species, *E. campestre*, called by John Ray, Friar's Goose.

THE SAMPHIRE.

(Fig. 41.)

CRITHMUM MARITIMUM.

THE SAMPHIRE is also an Umbelliferous plant. Those who have once seen it and smelt it will recog-

nize it again. It grows in places where none but the adventurous can reach,—on the sides of cliffs, near the sea, and in the clefts of rocks ; it fringes the edges of precipices with its bright-green succulent leaves and tiny blossoms. It is almost woody at the base ; the young branches, foliage, and umbels, thick and fleshy. The leaves are twice or thrice ternate, with thick linear segments about an inch long. The flowers are greenish, or yellowish-white, in umbels of fifteen to twenty or more rays. The samphire is warm and aromatic in flavour, and is frequently used as a pickle. Visitors to the sea-side who wish to try this pleasant condiment, cannot do better than look for it on the sides of the cliffs, and, if within reach, gather a basket-ful of its bright-green leaves ; they should be separated from the stalks and flowers, and then have spice and boiled vinegar poured on them in the usual way. Samphire-gathering, when pursued as a trade, frequently leads to loss of life, and terrible dangers are encountered to secure it. It has almost a classical association, since Shakspeare immortalized it in “King Lear.” It grows well on chalk rocks, such as form the cliffs at Dover ; there, Edgar is supposed to be leading Gloucester along, and says,—

“Come on, sir ; here’s the place : stand still. How fearful
And dizzy ’tis, to cast one’s eyes so low !
The crows and choughs, that wing the midway air,
Show scarce so gross as beetles. Halfway down
Hangs one that gathers samphire ; dreadful trade !
Methinks he seems no bigger than his head :
The fishermen that walk upon the beach
Appear like mice.”

It abounds also in the Isle of Wight, and many other sea-coasts of England, especially in the South, where, with its bright green colour, it gives quite a character to the cliffs and caves which overhang the sea.

WILD PARSNIP.

(Fig. 42.)

PASTINACA SATIVA.

THE WILD PARSNIP is very common on banks and in hedges when the soil is of chalk. It also belongs to the family Umbelliferæ. It has an angular, furrowed stem ; pinnate leaves, more or less downy, especially underneath, with five, seven, or nine segments, each one to three inches long, sharply toothed and more or less lobed. The flowers are yellow, in umbels, not very large, of eight to twelve rays. The root is spindle-shaped, and has the same appearance, odour, and taste as our well-known vegetable, which is, indeed, but a cultivated variety of the wild kind. In its native state, the root is tough and woody. It blooms during July and August, and its yellowish-green seeds remain on the stem until November. These contain an essential oil which once had a reputation as a cure for intermittent fevers. Hogs are fond of the roots, and quickly fatten on them. Parsnips have erroneously been supposed to contain a large quantity of nutritious matter. They are not so good for food as many other vegetables.

In the north of Ireland a kind of beer has been brewed from the roots, flavoured with hops; and the root yields by distillation an ardent spirit, from which a respectable sort of wine is made. In the colour of the flowers it is an exception to most of its family, as but few of the umbelliferous plants have any but white flowers

MISLETOE.

(Fig. 43.)

VISCUM ALBUM.

OUR next plant is associated with thoughts of pleasant meetings and festive boards, and happy are those whose homes are filled at Christmas time with the cheerful companions of summer rambles, having health and spirits to enjoy the good gifts of God in any form. The Mistletoe, or *Viscum album*, is one of a genus of parasitical plants belonging to the family Loranthaceæ, and is the only British representative of the family.

“Mistletoe,” says Lord Bacon, “chiefly grows on crab-trees, apple-trees, sometimes upon hazels, and rarely upon oaks, the mistletoe whereof is accounted very medicinal. It is an evergreen that bears a white glittering berry, and differs entirely from the tree whereon it grows. It continues green winter and summer, which the tree does not.”

“Nought was green upon the oak,
But moss and rarest mistletoe.”

In many respects the Mistletoe is an object of great interest to the naturalist. The manner in which it derives its nourishment from other plants by engrafting itself into the branches of a tree, and the curious conditions of the seeds in germination, are worthy of attention.

I need scarcely describe the plant so familiar to us all, as there is little fear of its being mistaken for any other. I may, however, say that the leaves are entire, varying from narrow oblong to nearly oval—thick, fleshy, and always obtuse. There are two kinds of flowers, those bearing pistils and those bearing stamens, each on separate plants: they are very small, and are placed in little heads, about three to five in each. The berry is about the size and has somewhat the appearance of a white currant, very smooth, viscid, and containing a simple seed. The manner in which the plant establishes itself in the branch of a tree has been much discussed. Old botanists believed that the “Mistletoe Thrush” feeding upon the berries surrounded his beak with the viscid mucus they contain, and in order to get rid of it, rubbed his beak against the branches, and thus inserted the seeds, from which springs a new plant. Paley, in his “Natural Theology,” gives at length his views on the subject. Of no other plant can it be said that the roots refuse to shoot in the ground, and no other plant is known to possess this adhesive generative quality when rubbed on the branches of trees.

The seeds in germination seem to offer an exception to the general law—that the radicle or root of the embryo shoots downwards, and the plumule upwards;

for it is found that the radicle of the Mistletoe invariably turns itself down upon the body to which it is attached, whatever may be the position of the surface of that body with respect to the earth. For instance, if a cannon-ball to which mistletoe seeds are glued on all sides be suspended by a cord some distance from the earth, both the upper and under seeds, as well as those on the sides, all direct their radicle to the surface of the ball. This property insures their growing upon the branches of trees, to whatever side they may happen to adhere.

It is asserted that a branch of Mistletoe, when placed in water, has no power of absorbing this fluid, but that when the branch to which it is attached is immersed, then the water is readily absorbed and penetrates into the Mistletoe itself. The following experiment was performed by De Candolle. He immersed the branch of an apple-tree bearing Mistletoe in water previously coloured with cochineal, which penetrating the wood and inner bark of the apple-tree, entered the Mistletoe, when its colour was even more intense than in the former.

Having no immediate connection with the earth, this curious plant, when discovered on the oak, already a sacred tree, became an object of superstitious worship to the priests of the Ancient Britons. The Druids, as they were called, held it in the greatest veneration, and the ceremony of separating the Mistletoe from the oak was one of their greatest religious rites. It was held on the sixth day of the moon, from which day they computed time. The occasion was celebrated by the sacrifice of two

white bulls, which were tied to the oak-tree by their horns ; then one of the Druids, clothed in white, mounted the tree, and with a knife of gold cut the Mistletoe, which was received in a "white sagum," or cloth made of wool : this done, they proceeded to their sacrifices and feastings. Our own national practice of decorating our churches and houses with mistletoe and holly at Christmas time may, perhaps, be a remnant of the old superstition. If, however, it only be regarded as a pleasant custom, there can be no harm in it, and long may it be ere so graceful a decoration be discontinued. Sir Walter Scott, in describing Christmas in the olden times, says—

"The hall was dress'd with holly green,
Forth to the wood did merry men go,
To gather in the mistletoe."

The origin of the old-fashioned tradition, that a kiss under the Mistletoe is only fair play, I am unable to find. I observe, however, as civilization advances, and manners become more polite, this practice is more honoured in the breach than the observance. In a pretty book I have had by me for many years, called "A Wreath of Friendship," written in part by the late Rev. Professor Henslow, when a young man, are some pretty lines on the Mistletoe. I cannot help quoting them here :—

"Past is the time, when bending low,
Druids revered thee, mistletoe ;
Error's broad shades are chased away
By Revelation's brilliant ray,
And superstition can no more
Bid us a humble plant adore.

Yet who in hour of Christmas mirth,
 Can place thee o'er the social hearth,
 With ivy and with holly gay,
 Or twine thee with the fragrant bay,
 Nor lift with joy his heart above,
 Nor hymn the notes of praise and love?
 Fair plant, a mystery thy birth,
 Thou dost not fix thy home on earth;
 Rock'd by the winds, fed by the shower,
 Thy cradle is an airy bower;
 No voice of crime in thy leafy dome,
 But the songs of birds to cheer thine home.
 From the wilding crab this branch was riven,
 From waving in the breath of heaven.
 Alas! alas! they have brought it low,
 To the dwellings of care, and pain, and woe."

 ELDER.

(Fig. 44.)

SAMBUCUS NIGRA.

THE COMMON or BLACK ELDER, is a small tree or large bush, belonging to the family Caprifoliaceæ. It is known by possessing a five-cleft calyx, a five-cleft rotate corolla, five stamens, and three stigmas, a roundish pulpy one-celled berry, hardly crowned by the remains of the calyx. The stem is irregularly but always oppositely branched, the young branches are clothed with a smooth grey bark, and filled with a light spongy pith; the leaflets are deep-green and smooth, usually with an odd one. The flowers are in bunches of a cream-colour, with a sweetish but faint smell. The berries are black, with reddish stalks.

Considerable medicinal value was at one time attributed to this plant, but it is now chiefly used in the rural districts of England for making a wine from the berries, which, when spiced and drunk hot, is regarded by some people as very agreeable. The flowers are employed in making a distilled water, which is cooling and refreshing, and is sometimes introduced into confectionery. The wood is hard and tough. The pith, on account of its solidity and great lightness, is used for making small figures and balls for electrical experiments. The undeveloped buds when pickled, form a good substitute for capers. It is on the leaves of the elder that the caterpillar of the sphinx or death's-head moth delights to feed. It changes into a chrysalis about September, and in the following July may be seen as the gigantic and curious moth, with the markings of a skull on its thorax. There are two species of elder native in Britain, of which *S. Ebulus*, or Dwarf Elder, a rare plant, is the other.

HONEYSUCKLE, OR WOODBINE.

(Fig. 45.)

LONICERA PERICLYMENUM.

THIS is a favourite flower, on account of its delicious fragrance and its lovely blossoms, intermingling with every hedge-row, and turning round the trunk of many a sturdy tree. It belongs to the family Caprifoliaceæ. The leaves are ovate or oblong, smooth

above, somewhat downy or slightly hairy beneath,—the upper ones the smallest. The flowers are of a pale-yellow colour, the corolla about an inch and a half long. The berries are small and red. There are two other species of honeysuckle native in Britain,—*L. caprifolium*, the Perfoliate Honeysuckle, or Goat's-leaf, and *L. Xylosteum*, the Fly or Upright Honeysuckle.

The leaves of the perfoliate honeysuckle are quite smooth and broader than those of our example, and it flowers earlier in the year. The fly honeysuckle is rather rare; it has white scentless flowers, and is by no means so attractive as either of the others. It is often planted in shrubberies.

The true Woodbine of the poets is undoubtedly the *L. Periclymenum*; it obtains the name evidently from a corruption of *woodbind*, from its habit of twisting round the stems of trees. Milton calls it “the Twisted Eglantine,” and Shakspeare says—

“So doth the woodbine, the sweet honeysuckle,
Gently entwist the maple.”

At the base of its long tubular flower lies the honey, and when the bee cannot reach it, other insects tap it, by making a puncture at the base of the tube, and thus regale themselves. In almost every country lane in England, from early in June to August, are we delighted with the sweet scent of one or other of these pretty climbers; and well do I remember, on a memorable occasion, becoming almost overpowered with the perfume, as bough after bough was pulled from its native Suffolk hedge, and piled up, so as to

make a floral couch in a carriage in which I sat one still warm summer evening in July. Later in the year, the clusters of bright red berries, which follow the flowers, are very picturesque, and afford food for the birds; for they are not poisonous. The hawk-moth is often found hovering near the honeysuckle, attracted, perhaps, by its fragrance or feeding on the honey it contains.

YELLOW BEDSTRAW.

(Fig. 46.)

GALIUM VERUM.

LADIES' BEDSTRAW, or Cheese Rennet, belongs to the family Rubiaceæ. It is distinguished by having its leaves about eight in a whorl, small, and linear. The stems are much branched at the base, six inches to a foot long, ending in a panicle of small golden-coloured flowers. The roots afford a rich yellow dye, and impart this colour to the bones of animals who feed upon it. The whole genus possesses a property like that of rennet, of curdling milk, which gives rise to their common name. The *Galium verum* is one of the prettiest plants, which decorates our driest sand-banks, gaily blossoming for full three quarters of the year. It is the sweetest of all the genus, and was formerly much used for strewing floors and laying in beds; whence the name of bedstraw, at a time when feather-beds and luxurious spring couches were unknown. According to John Ray, the flowering tops, when distilled, make a refreshing beverage,

and the roots are useful as an astringent medicine. The French used formerly to prescribe the flowers in hysteria and epilepsy.

FULLER'S TEAZLE.

(Fig. 47.)

DIPSACUS FULLONUM.

THE FULLER'S TEAZLE will be easily recognized, from the drawing, although we have one or two other British species. It belongs to the family Dipsacæ, and is a stout biennial, four or five feet high, with numerous prickles on the stems, the leaves, and, in fact, on the whole of its surface. The heads of the flowers are of a pale lilac-colour, at first ovoid, but gradually becoming cylindrical, nearly three inches long and about one inch and a half in diameter; the scales of the involucre are hooked, or reflexed, and very hard. On this account they are used in the manufacture of cloth in a process of brushing or raising, called *teazling*. It consists in applying the ripened head or fruit of the teazle to the cloth. The teazles are attached to a cylinder, which revolves upon the cloth, and the loose particles are raised, so that they may be easily sheared or cut off, to give the cloth the fine appearance it assumes. No instrument has ever been invented to supersede the teazle. Various substitutes have been tried; pieces of wire have been fixed into a leather back, but nothing answers the purpose so well as the elastic spines of the teazle.*

* See Dr. Lankester's Lectures on the Uses of Animals.

The plant is imported in large quantities from France and other parts of Europe, and is also extensively grown in England ; but as a hot sun and dry weather are essential to the proper drying of the teazle heads, those imported from the Continent are esteemed the best. Country people are still said to cure agues in various parts of England by a singular remedy obtained from this plant, which is of course only imaginary. If the heads be opened longitudinally in the autumn, a small worm may frequently be found in the centre. Of these, three, five or seven, always an odd number, must be taken, sealed up in a quill and worn in good faith, as an amulet or charm against ague.

CHICORY.

(Fig. 48.)

CICHORIUM INTYBUS.

CHICORY, or Succory, belongs to the family Compositæ. It is often seen growing wild on the borders of our corn-fields, and is sure to attract attention by its pretty blue flowers. It is from one to three feet high. The leaves near the ground are spreading, more or less hairy, with a large terminal lobe and several smaller ones, all pointed and coarsely toothed, the upper leaves smaller, less cut, and embracing the stem by pointed auricles. The flowers are in heads, in closely sessile or unstalked clusters, of a bright blue colour. When blanched, or grown without light, the

49



50



51



52



53



54



55



56



57



58



59



60



leaves are often eaten in early spring salads, and are very good ; they are somewhat bitter, like endive, but lose that taste by cultivation. The French call the long slender leaves "Barbe de Capuçin," Monk's Beard. The root is long, like a carrot, and is used in large quantities, when dried, as a substitute for, or an addition to, coffee. In Belgium and many parts of Germany, large districts are devoted to the cultivation of this plant for the sake of its root, which is dried in a kiln or slow oven. It is afterwards roasted like coffee, ground in a mill, and sold in the market. It is decidedly a pleasant addition to coffee ; but, being much cheaper than the coffee berry, it becomes an adulteration when mixed with it and sold as pure coffee. If a teaspoonful of the powdered chicory be added to a teacupful of ground coffee, it will be found to improve the flavour, and to be in no way injurious.

THE SEA ASTER.

(Fig. 49.)

ASTER TRIPOLIUM.

THE SEA ASTER, or Sea Starwort, is a pretty plant, found growing in salt marshes, near the coast, and belonging to the family Compositæ. It is seldom above a foot high, erect or decumbent at the base ; slightly branched, the leaves linear, entire, and somewhat succulent. The flower-heads are in a compact corymb ; the involucral bracts few and oblong. The centre of the disk of the flower is bright-yellow, the circumference blue or purple.

ELECAMPANE.

(Fig. 50.)

INULA HELENIUM.

ELECAMPANE is found in pastures in various parts of Europe. It is a native of Great Britain, and belongs to the order Compositæ. It has a thick, branching root, which is aromatic, bitter, and mucilaginous. The stem is three feet high, leafy, round, furrowed, solid branched, and most downy in the upper part. The leaves are large, ovate, serrated, and veiny, downy and hoary at the back; the root-leaves stalked; the rest are sessile, clasping the stem. The flower-heads are solitary, at the downy summits of the branches, two inches broad, of a bright-yellow colour, with reddish streaks; the scales of the involucre are broad, recurved, leafy, finely downy on both sides. The rays are very numerous, long, and narrow, each ending in three unequal teeth. Various preparations of the boiled root have been recommended, mixed with sugar, to promote expectoration and to strengthen the stomach. Some think a spirituous extract contains most of its aromatic and tonic properties. We may generally find the plant in cottage gardens, on account of its reputed virtues. Inulin, a peculiar substance contained in the root, is a form of starch insoluble in cold water, but soluble in hot water, from which it is deposited on cooling. With iodine it gives a greenish-yellow compound which is not permanent.

Inulin is distinguished from gum by its insolubility in cold water, and is otherwise chemically interesting.

CHAMOMILE.

(Fig. 51.)

ANTHEMIS NOBILIS.

COMMON CAMOMILE, or Chamomile, is frequently found in a wild state on many of the commons near London, where it adds a peculiar richness of colour and fragrance to the turf. It is a dwarf plant, belonging to the family Compositæ, with finely-cut leaves; the flower-heads are white in the ray, but deep-yellow in the disk. All parts of the plant are intensely bitter, especially the little yellow flowers of the disk; for this reason the wild flowers are more efficacious than the cultivated sort, in which there is scarcely any disk, the white flowers of the ray having almost entirely usurped their place. An infusion or extract of these flowers is often used in medicine as a stomachic, and also as a fomentation externally. Besides the bitter principle contained in this plant, chemists have obtained from it camphor and tannin, and also a volatile oil, of a beautiful blue colour.

HAREBELL.

(Fig. 52.)

CAMPANULA ROTUNDIFOLIA.

No wild flower is more admired, or has had its praises sung by poets more frequently, than this pretty

delicate little inhabitant of every heath and sunny bank of our country districts. Every village child loves its pretty bells, and numberless are the fancies which connect it with fairy legends and floral charms. It seems scarcely necessary to describe it botanically; but, lest it should be confounded with other species of the same genus, it may be well to say that it belongs to the family Campanulaceæ. The leaves on the tiny stem are very slender, like those of grass, but near the ground there are a number of roundish notched leaves, which mostly die away at the time of flowering. The bell-shaped corolla is of a pale blue colour, and has five broad lobes, much shorter than the entire fruit. We have nine wild species of *Campanula*, some of which have stout stems with large purple flowers, many of which bear the cultivation of the garden very much better than our true little Harebell, which is unhappy away from its native haunts. The Canterbury Bell, with its large handsome flowers, is one of the favourites of our gardens, and at one time abounded in the neighbourhood of Canterbury and other parts of Kent: it was gathered by pilgrims to the shrine of St. Thomas à Becket there, and treasured in evidence of the task they had completed. The little plant I have chosen as most worth notice is the true Harebell of Scotland, the same which in the "Lady of the Lake" is mentioned as being strewn in Ellen's pathway.

"For me she stoop'd, and looking round,
Pluck'd a blue harebell from the ground;
This little flower that loves the lea
May well my simple emblem be."

It is said that the presence of the Harebell indicates a barren soil ; yet how lovely are its tiny cups on their cobweb stems, gently waving to and fro with every breath of wind, so that one might almost believe in the reality of the silver music said to come from them in the days of yore, when the good fairies

“Rang their wildering chimes to vagrant butterflies.”

And even now, with all the sobering influences of botanical study upon us, we can heartily sympathize with the little one who, having filled her lap and both hands with blue bells innumerable, and *white* bells, too, both growing close together, was heard to whisper in the real spirit of prayer and happiness—
“Dear God, do make some pink bells too !”

BLACK WHORTLEBERRY.

(Fig. 53.)

VACCINIUM MYRTILLUS.

THIS plant, known also as Black Whortleberry, may be regarded as the representative of the British berries known as Bilberries, Crauberrys, Cowberries, Windberries, &c. It belongs to the natural order or family Ericaceæ, and is a small shrub, from six inches to a foot high, with spreading green branches. The leaves fall off in the winter, and are small, ovate, with tiny teeth, and a very small stalk. The flowers are nearly round, of a pale greenish white colour, with a tinge of red, growing singly in the axils

of the leaves. The berries are round, nearly black, and covered with a bluish kind of bloom, crowned by the short teeth of the calyx. This shrub is found on most mountain heaths and woods throughout England, Scotland, and Ireland, with the exception of the eastern part of England. The leaves have been much used in the adulteration of tea. The berries are frequently eaten, either in tarts or uncooked. They have a sharp astringent flavour, which is not pleasant to every one, but, like the rest of the family, they have their admirers.

The Cranberry, *V. Oxycoccus*, is well known, and is much liked in England. Great quantities of the berries are imported from Russia, Sweden, and America, into this country, packed in tubs. They are considerably larger and finer in appearance than those grown in our own country, but not so full of flavour.

The other species of *Vaccinium* are the Bog Whortleberry, *V. uliginosum*, and the Red Whortleberry or Cowberry, *V. Vitis-idea*.

HEATH.

(Fig. 54.)

ERICA TETRALIX.

THIS is the most widely distributed and best known of all our native heaths. The leaves are four in a whorl, lanceolate and linear, ciliate, downy above and on the midrib beneath. The stem is bushy at the base, with rather short, erect, flowering branches.

The flowers are rose-coloured, forming little clusters or close umbels at the end of the stalks. It is commonly found in the West of Europe, and in Britain is most plentiful in the western counties. I have, however, constantly found it in Suffolk, the Isle of Wight, and other parts, often growing with the heather or *Calluna vulgaris*, from which it is well to distinguish it. The larger and more bell-like blossoms of the *Erica*, and its downy appearance, are the evident marks by which we may recognize it. These heath flowers were adopted as the the badges of the Highland clans; and although this heath is not especially a Scotch plant, the *Erica tetralix* was borne by the Macdonalds, the *Erica cinerea* by the Macalisters, and the *Calluna vulgaris* by the Macdonnells. All these plants grow together on the moors and fells in the North of England and Scotland, and give a peculiar aspect to the landscape, shedding as it were a purple hue over the distant mountains, and forming a characteristic feature of these northern districts.

There are six species of *Erica* in Great Britain.

HEATHER.

(Fig. 55.)

CALLUNA VULGARIS.

THE HEATHER grows, as I have said, in common with the heaths, and is often mistaken for them. Its smaller more purple blossoms, placed all along the stems in little bunches, will serve to distinguish it.

Sometimes the flowers are white, but this is rare. It is especially the plant of the Highlander, and is associated so strongly with his country in all its legends and poetry, that it appears almost as exclusively the child of the mountain fastnesses as the national music of the bagpipe. To the Highlander this plant is something more than a mere badge of clanship; it furnishes him with much that is valuable in everyday life. The heather branches, freshly gathered and arranged so that the elastic tips of the shoots form a level surface, constitute a couch such as that described by Sir Walter Scott:—

“ Before the heath had lost the dew,
This morn a couch was pull’d for you,
On yonder mountain’s purple head.”

And again,—

“ The stranger’s bed
Was there of mountain heather spread.”

Cabins are also thatched with it, and the walls of the cottages are often made of alternate layers of heather and a kind of mortar. As fuel it serves well, and it is said to yield a yellow dye, which I am told is at present used by the cloth-manufacturers of Yorkshire. Moreover, in England the sprigs of the heather are constantly made into brooms or besoms, which are very serviceable. As food for moor game and grouse, the heather is almost essential, and it is only where this plant will grow that these birds can be preserved. The red deer also crops the young shoots of the heather. Bees extract honey from the flowers, which, though dark in colour, is very rich in flavour.

I have seen the heather, and, indeed, many species of heath, prettily used as a border for flowerbeds in gardens. Sir W. Hooker suggested it, and has carried it out at Kew. Accustomed as we are in the southern districts to see the heath plants only as a low shrub, a foot or two in height, we are surprised to read of—

“Heather black that waved so high,
It held the copse in rivalry.”

Yet so it is, and in certain wild and peaty districts it may be found quite tall enough to justify this description.

THE HOLLY.

(Fig. 56.)

ILEX AQUIFOLIUM.

THIS plant belongs to the small natural family Aquifoliaceæ, and is the only British representative of the family. It seems hardly needful to describe so well-known and favourite a tree, associated as it is with the happiest days of childhood, with Christmas-day gatherings and merry-makings, with joyous faces and warm hearts, while to some of us, perhaps, who have passed the sunshine of life, its bright green leaves and red berries may call up memories of the companions of past years, now passed away, never more to share in our joys or our sorrows. Familiar as we all are with the red berries of the Holly, we may not have seen its flowers, for they blossom when

all nature is bright, and are overlooked amidst their more showy and attractive neighbours. They grow closely round the stem, and are white and wax-like, opening in May and June. The bright shining green leaves are armed with sharp prickly teeth, but the upper ones on a bush are frequently smooth. This circumstance the poet Southey impresses on the memory in his charming lines on the Holly Tree,—

“Below a circling fence its leaves are seen,
 Wrinkled and keen.
 No grazing cattle through their prickly round
 Can reach to wound ;
 But as they grow where nothing is to fear,
 Smooth and unarm'd the pointless leaves appear.”

The tendency to produce these prickly points renders the Holly peculiarly fit for hedges, and when Dutch horticulture prevailed in England, such hedges were not unfrequent.

“A hedge of holly, thieves that would invade,
 Repulses like a growing palisade.”

The celebrated John Evelyn had such a hedge at Say's Court, four hundred feet long, nine feet high, and five feet broad, which he planted at the suggestion of Peter the Great, who resided at his house while he worked in the Deptford Dockyard. In his Diary he asks, “Is there under heaven a more glorious and refreshing sight of the kind than such an impregnable hedge, glittering with its armed and varnished leaves, the taller standards at orderly distances blushing with their natural coral?” The Holly is a very slow-growing tree, and its timber is

amongst the hardest of white woods ; it is much used by turners, and especially in the manufacture of Tunbridge ware.

THE SPRING GENTIAN.

(Fig. 57.)

GENTIANA VERNA.

THIS is one of the brightest ornaments of our northern districts. It belongs to the family Gentianaceæ, and to a genus which flourishes especially in Alpine and Arctic regions. This pretty little Gentian is, however, sometimes found in warm mild districts. I have met with it in the Isle of Wight, on Shanklin Downs. Bitterness is a characteristic of the whole family, and this principle seems to have something to do with the bright blue colour of the blossoms, for the brighter the hue the more bitter the taste. Our little spring Gentian has a perennial leafy stock, densely tufted, often spreading to four or five inches in diameter, with ovate or oblong leaves. The flower-stems are simple and numerous, sometimes so short as to give the flowers the appearance of being seated on the leaves ; sometimes they are an inch or two in length, and bearing one bright-blue terminal flower. The corolla is tube-like, nearly an inch long, with five ovate lobes, and smaller two-cleft ones between them. The bitter principle of all the gentians is valuable as a medicinal agent.

G. nivalis, the little Snow Gentian, grows on our loftiest mountains in Scotland and Wales, but is far better known as a native of the Alps and Pyrenees.

BUCKBEAN, OR MARSH TREFOIL

(Fig. 58.)

MENYANTHES TRIFOLIATA.

THE BUCKBEAN belongs also to the family of Gentians—Gentianaceæ. It is a beautiful aquatic herb, and is found in wet bogs and shallow ponds all over Great Britain. The stem is short, creeping, or floating, with a dense tuft of leaves, consisting each of a long stalk, sheathing at the base, and three obovate or oblong leaflets, one to one and a half inches long. The flowers are white, tinged externally with pink, in an oblong raceme on a peduncle of from six inches to a foot long, proceeding from the base of the tuft of leaves. The corolla is deeply five-lobed, and fringed on the inside with white filaments. None of our native plants exceed this in beauty. In the fresh-water aquarium it is a beautiful object, and may easily be preserved in this artificial condition for some time. In common with all its family, the Marsh Trefoil abounds in an intensely bitter quality, which has frequently been used medicinally. Withering says that the leaves have been used during a scarcity of hops as a substitute for them in brewing beer.

85



Yellow Orchid

86



Daffodil

87



Wild Hyacinth

89




Ribwort's Snail's shell

88



Grape Hyacinth

90



Snakeshead Lily

91



Turkey Tulip

92



Meadow Saffron

93



Flowering Rush

94



Arrow-head

95



Cuckoo-pint

96



Sweet Flag

GREAT BINDWEED.

(Fig. 59.)

CONVOLVULUS SEPIUM.

THERE can scarcely be any occasion to describe this favourite and well-known plant, which wreathes its graceful festoons over our hedgerows, opening its large tender white blossoms to the bright sunshine, and gathering their folds together as a rain-cloud foretells the approach of a shower, which would shatter their delicate texture. Botanists no longer call this plant by the old name so familiar to our childhood. It is no longer the *Convolvulus*, but is known as a distinct genus, under the name *Calystegia*. The lovely white blossoms survive but for a single day; whence they are called "Belle de jour" by the French. They are, however, so rapidly succeeded by a profusion of buds ready to take their places, that the decay is not noticed, and our attention is taken off from the flower which has lived "its little day" and is now no more. A well-known author says, "How affecting an emblem of human life does this simple *Convolvulus* present to us. The gay, the young, whose existence has seemed but a day, are cut off, and others, equally gay and equally mortal, occupy their places; and the remembrance of them is quickly dissipated by the attractions of their successors, who, perhaps, like them, are doomed early to submit to the common lot of humanity."

Beauty alone is not the sole merit of this plant ; the root has properties similar to those of *C. Scammonia*, and has been used as its substitute under the names of Montpellier and Bourbon Scammony. It has an ancient reputation as a medicinal agent, Galen himself being said to have recommended the leaves as an external application to swellings and abscesses.

SEA BINDWEED.

(Fig. 60.)

CONVOLVULUS SOLDANELLA.

THIS is one of the prettiest of our sea-shore plants. It is abundant on the eastern coasts of England, on sand-hills, and flourishes well on the red clay of Suffolk and Norfolk. It belongs to the same family as the large *Convolvulus*, but is much smaller, trailing along the ground, and seldom rising to a height above six or eight inches. The leaves are small and kidney-shaped, with broad lobes at the base. The flower is of a delicate or bright pink colour, streaked sometimes with a darker shade. This plant shares, in common with the rest of its family, a reputation for curative properties ; indeed, but few of our wild plants have not been employed in rustic pharmacy with more or less success ; and it is only since the science of chemistry has proved that the active principles of these vegetable productions are the same in many different plants, and can be extracted by chemical processes, that the practice of making infu-

sions of all sorts of herbs and weeds has been gradually abandoned for a more convenient method of administering the valuable principles to be found in nature's laboratory in smaller quantities and a less troublesome form.

VIPER'S BUGLOSS.

(Fig. 61.)

ECHIU M VULGARE.

THIS plant is one of the most beautiful of our wild-flowers. It belongs to the family Boraginaceæ, and is known by its showy flowers, having an irregular and unequal margin, and a sort of bell-shaped figure. The red or purple spots and hairs are very remarkable. The corolla is at first of a reddish-purple colour, turning afterwards bright-blue; so that there are constantly flowers of both colours to be seen at once. The whole plant is covered with stiff, spreading, almost prickly hairs. The root-leaves are stalked and spreading, but often withered away at the time of flowering. The common name of Viper's Grass originated from a resemblance the ripe seeds are supposed to have to the head of that reptile; and hence arose the idea that it might act as a remedy against the bite of the creature,—an adaptation of the more recent but not less absurd fallacy that "like cures like." The showy blossoms are very attractive to bees, and not even the sharp hairs by which they are guarded are sufficient to deter these little plunderers

from their depredations on the sweet store concealed in the flowers. But we must not consider these little creatures as merely selfish seekers of their own gratification ; they, in common with the whole of creation, serve great purposes, and carry out the designs of the Great Architect of all, even without their own consciousness of doing so. As the insect flies from flower to flower, it transports on its delicate legs and wings some portion of the pollen which is to fertilize the next plant which it enters, and produce seed to reproduce the species. Every class of animals seems to assist in this great work of propagation in the vegetable world. Seeds are carried about by birds and deposited, as if by accident, in situations favourable for their growth ; and even the giddy butterfly as it alights on each gay flower in the sunshine, is the messenger of a new life to the rapidly-fading beauty of the garden.

FORGET-ME-NOT.

(Fig. 62.)

MYOSOTIS PALUSTRIS.

THIS is so entirely a flower of associations, that it is difficult to unravel anything like a botanical or prosaic description of the plant itself, from the numerous poetical fancies and legendary tales by which it is surrounded. It belongs to the Borage family,—Boraginaceæ, and has small downy, ovate leaves, which are not unaptly compared to mouse-ears. Botanists, finding that the plants commonly received as Forget-

me-nots differ in some minor characteristics, divide the species into three varieties. The true Forget-me-not has rather a large rotate flower, of a clear blue colour, with a yellow eye. Most abundantly does it grow beside brooks, rivers, and wayside streams, and—must we say it?—even in stagnant ditches, asking only for moisture to adorn the most deserted places with its turquoise flowers. In cultivation, it will even dispense with this requirement, and will produce blossoms of a larger size than when wild. It is an excellent plant for window gardening, and is improved by “bedding,” as the gardeners call it—blossoming all the summer through, if properly trimmed. Beauty, however, is not the sole attraction of this favourite flower; it has associations connected with it in legends, in poetry, and in real life, which live long after its beautiful blossoms have perished. For many centuries it has been regarded throughout Europe as the emblem of eternal friendship or love. It is pleasant to regard so lovely a flower as expressive of a tender feeling. The well-known story is one belonging to the days of chivalry, when a knight and his lady-love were wandering on the banks of a stream where grew clusters of these gem-like flowers. In those days the wish of a loved one was law to the lord: the lady, desiring to possess some of the bright blue blossoms, caused her faithful knight to rush into the stream to obtain them for her, when, overborne by the strength of the current, he was carried away, and could but cast, with dying hand, the flowers she wished for towards her, exclaiming “Forget me not.” Even in our less chivalrous and more prosaic times, the language of this

flower is not forgotten, and it is not long since we saw, on St. Valentine's day, when such sentiments are in fashion, and all nature is pressed into love's service, a pretty painted wreath of "Forget-me-nots" inclosed in a suitable envelope, with the words "Pretty flowers, speak for me," neatly inscribed within. One of our great botanists suggests, with a more philosophical and less poetical mind, that the real signification of the name is, after all, due to the bright blue tint and yellow eye of this charming flower, which, if once seen, is not likely to be forgotten. Nevertheless, we remain faithful to the generally-received tradition, and in justification of it, call to remembrance that as early as 1465, when a joust was held in which Lord Scales, brother to the Queen of Edward IV., took part, the fair ladies of her court presented to that favoured knight a collar of gold, enamelled with "Forget-me-nots." I am not aware that this much-prized plant has ever been used in the arts of life, yet it is a household favourite, and reminds us that there is in the human mind a deep and close association between the external beauty of nature and the strongest feelings of our hearts. Who but loves to meet, as Coleridge has it,—

"By rivulet or wet roadside,
That blue and bright-eyed flow'ret of the brook,
Hope's gentle gem, the sweet Forget-me-not."

HENBANE, OR HOGSBEAN.

(Fig. 63.)

HYOSCYAMUS NIGER.

THIS belongs to the family Solanaceæ, which is chiefly characterized by its poisonous properties. Wherever there is a patch of waste ground, there may be seen the dull-yellow blossoms of this dangerous plant. Its whole appearance, and the peculiar faint and disagreeable odour emitted by it, would seem almost to indicate its nature; yet the narcotic principle yielded by it is most valuable as a medicine, properly administered. It is a coarse, erect, branching annual, about one or two feet high, more or less hairy and viscid. The leaves are rather large, sessile; the upper ones clasping. The calyx is short when in flower, but points round the fruit, and is then an inch long, strongly veined, with five broad, stiff, almost prickly lobes; the stem ovate, and irregularly pinatifid. The flowers are remarkable for their purple veins, which give them a curious appearance when in blossom, and which cause the whole plant to be peculiarly adapted for a process now not uncommon in ornamentation. If soaked for a long time in water, with some chemical agent, the soft parts of a plant decompose and disappear, leaving only the woody fibrous parts entire and bleached. These form what are known as skeleton plants, and I have seen several extremely beautiful specimens of the Henbane treated in this way, which indicates that the

woody tissue is abundant in this plant. The narcotic properties of the Henbane are most strongly developed when the flowers have just fallen and the seeds are ripening. Lightfoot mentions that a few of these seeds have deprived a man of his reason and the use of his limbs; nevertheless they are frequently smoked in a tobacco-pipe as a cure for toothache. The expressed juice of the plant, forming an extract, is recognized in the "Pharmacopœia," and in some cases is very valuable where other anodynes are inadmissible. A curious effect is produced on the system by these vegetable poisons, somewhat like intoxication—excitement, restlessness, and irritability are often the first symptoms of their action. A story is told of a gardener and his wife, who lived happily and in perfect contentment, until one day the good man, wishing to dry some Henbane plants, hung them up in his bedroom for that purpose. From that hour his domestic peace vanished; his wife became a perfect shrew, and he returned each curtain-lecture with interest. Happily Sir Cresswell Cresswell's court was then unknown, or the speedy separation of the discontented parties would have rendered the solution of the mystery for ever impossible. Accidentally the Henbane was removed, and peace was restored. Each felt that, after all, the other was not to blame, and with returning amiability came increased happiness. It remained, however, for philosophers to trace the connection between the baneful effects of the Henbane exhalations and the irritable, quarrelsome condition of those who breathed them.

DEADLY NIGHTSHADE, OR DWALE

(Fig. 64.)

ATROPA BELLADONNA.

THIS is one of our most poisonous native plants, and must be clearly described in order to be avoided. It belongs to the family Solanaceæ, and is an erect, smooth, or rather downy herb, with a perennial root, stock, and branching stem. The leaves are stalked, rather large, ovate, and entire, with a smaller one usually proceeding from the same point, often so small as to look like a stipule. The flowers are solitary, on short peduncles in the forks of the stem, or the axils of the leaves. The flower is of a pale purplish-brown colour, bell-shaped, nearly an inch long. The stamens are shorter, with distinct filaments. The berries are large, and of a shining black appearance, which frequently tempts children into eating them, to their injury. The modern name *Belladonna*, refers to the practice of the Italian *belles*, who make use of its properties to enhance their personal charms. A portion of the extract, when placed in contact with the pupil of the eye, causes it to dilate, and gives a brilliancy and lustre to these speaking orbs, which is much coveted and admired. I am told that this practice is not confined to the land of cloudless skies and southern breezes, but that in our own country the preparation is to be seen on the toilette-tables of our fashionable ladies. Happily this property is

turned to good account by modern science, and in examinations of the eye it is found to be of great service in dilating the pupil, as well as previous to the operation for cataract. Numberless are the instances where death has ensued from partaking of this plant or its berries. The very powerful nature of its poisonous qualities has directed the attention of modern professors of *Materia Medica* towards it, and at this time it is considered to be a valuable medicinal agent. A medical friend of ours, enthusiastic in the discovery of hidden truth, not long ago nearly fell a victim to his own experiments on the action of an extract of the leaves of *Atropa Belladonna*. He had previously made known his belief that animal charcoal is the best and safest antidote to vegetable poisons. He was, however, too much under the influence of the poison to think of his own remedy, and it was only by the timely interference of a friend, who knew of the discovery, that his life was saved. It is a favourite remedy in homœopathic medicine; but as the doses given are inappreciable, it would be difficult to trace any results from their administration. The poisonous properties of this plant have long been known, as appears from its having been used by the Scotch under Macbeth to poison the Danes. Our great poet Shakspeare, with his wonderful appreciation of natural phenomena, refers, undoubtedly, to the same plant in Banquo's speech,—“Or have we eaten of the insane root that takes the reason prisoner?” Paroxysms of madness are among the curious and direful effects of this plant on the system, to which it is supposed Plutarch refers in his account of the strange

and disastrous results produced on Marc Antony's soldiers, "from tasting unknown herbs" when distressed for provisions.

To the same genus of plants belong the Mandrakes, *Atropa mandragora*, the roots of which are superstitiously connected with numerous fancies, and are still sold on the continent of Europe as ingredients in love-philtres and charms. Some writers recognize them as the mandrakes of Scripture.

WOODY NIGHTSHADE, OR
BITTER-SWEET.

(Fig. 65.)

SOLANUM DULCAMARA.

THIS belongs to the same family of plants as our last example, and is believed to have the same poisonous properties, in both fruit, leaves, and stem. This has, however, been disputed of late, and the Professor of Materia Medica at University College considers all the Solanums as perfectly innocuous. It has a shrubby stem, with climbing or straggling branches, often many feet in length, but dying far back in the winter. The leaves are stalked, ovate, or lanceolate, two or three inches long, usually broadly cordate at the base and entire, but sometimes with an additional smaller lobe or segment on each side, either quite smooth or downy on both sides, as well as the stem. The flowers are rather small, blue, with yellow

anthers, in loose branches, shorter than the leaves. The berries are small, round, or ovoid, and of a bright-red colour. It is to be found in hedges in shady, moist situations, all over England and Ireland—more rarely in Scotland. As a medicine, this plant has been used both internally and externally; it is recommended in asthma and many other diseases.

GERMANDER SPEEDWELL.

(Fig. 66.)

VERONICA CHAMÆDRYS.

THIS is one of the prettiest of our wayside plants, belonging to the family Scrophulariaceæ. There are so many species of *Veronica*, that they are somewhat difficult to distinguish from each other. The lovely blue flower, "The Celestial Bird's-eye Blossom," as it has been called sometimes, causes it to be mistaken for the true "Forget-me-not," but those who study both will soon learn to distinguish them. The leaves are shortly stalked, ovate, cordate, crenate, and hairy. The flowers spring from the leaves on rather long stalks, and have a five-cleft calyx. The common name of Speedwell is well chosen, for what so cheering to the wayfarer as its bright-blue flowers peeping out from the hedgerows to greet him as he passes by—"Stars that in earth's firmament do shine." The whole of the *Veronicas* have had their day in rustic medicine, and perhaps a larger list of virtues belongs to the

Germander Speedwell than any of the rest. As a substitute for tea it is considered excellent by some writers. The Emperor Charles V. is said to have used it as a remedy for gout, and in cancer it is recommended by old Gerarde to be given in "Good broth of a hen," a prescription in which possibly the *vehicle* may have had more efficacy than the medicine.

In a pleasant little book by Mr. Hibberd, called "Brambles and Bayleaves," an anecdote connected with this charming little plant is given from the life of Rousseau, which is too interesting to remain unquoted here. "During the earliest and happiest days of Rousseau's life, he was walking with a beloved friend on a calm, serene, summer evening. The sun was setting in all its glory, spreading sheets of fire over the western sky and upon the unrippled surface of the lake, making the water still more transparent with a vivid and glowing light. The friends sat on a soft mossy bank, enjoying the calm loveliness of the scene, conversing on the varied phases of human life in the unaffected sincerity of true friendship. At their feet was a bright tuft of the lovely Germander Speedwell, covered with a profusion of brilliant blue blossoms. Rousseau's friend pointed to the little flower, the *Veronica Chamædrys*, as wearing the same expression of cheerfulness and innocence as the scene before them. Thirty years passed away! Careworn, persecuted, disappointed, acquainted with poverty and grief; known to fame, but a stranger to peace, Rousseau again visited Geneva. On such a calm and lovely evening as thirty years before, he had con-

versed with the friend of his bosom, and had received a lesson from the simple beauty of a flower, he again was seated on the selfsame spot.

“The scene was the same! The sun went down as before in golden majesty; the birds sang cheerfully in the soft light of eventide; the crimson clouds floated solemnly in the western sky, and the waters of the lake were calm as before. But the house wherein the first feelings of love and friendship, and the first fruits of his genius had budded, was now levelled with the ground. His dearest friend was sleeping in the grave. The generation of villagers, who had partaken of the bounty of the same beneficent hand, had passed away, and none remained to point out the green sod where that benefactor lay. He walked on pensively; the same bank, tufted with the same knot of bright-eyed speedwell caught his eye, the memories of past years of trouble and sorrow came upon him; he heaved a sigh and turned away, weeping bitterly.”

SNAPDRAGON.

(Fig. 67.)

ANTIRRHINUM MAJUS.

THIS is a plant belonging also to the family Scrophulariceæ. It is frequently cultivated in gardens, but also grows wild on old walls and stony places, and on chalk cliffs on the south of England. The leaves are narrow, lanceolate, or linear entire. The flowers are large, of a purplish-red colour, or variegated with

white. The corolla is above an inch long, opening like a mouth when pressed between the finger and thumb. They form perfect insect-traps : numbers of these little creatures, attracted by the sweet nectar to be found within, enter the treacherous tube, but on seeking to return, find their egress effectually prevented. Those, however, who are clever enough to gnaw a hole through the side of their prison escape uninjured. Gmelin says, that in Persia an excellent oil, equal to that of the olive, is procured from the seeds of the *A. majus* ; and Vogel observes, that the common people in many countries attribute some supernatural influence to this plant, believing it to have the power of destroying charms, and rendering maledictions of none effect. All the varieties of Snapdragon are peculiarly able to resist the effect of great droughts, and supply vegetation and beauty in situations where other flowers would perish from the power of the sun's rays.

GREAT MULLEIN.

(Fig. 68.)

VERBASCUM THAPSUS.

THIS plant belongs to the natural family Scrophulariaceæ, and is a stout, erect biennial, two to four feet high, clothed with soft woolly hairs, which circumstance I believe has given rise to its name *Verbascum*, being a corruption of *barbascum*, or bearded. The leaves are oblong, pointed, slightly toothed, nar-

rowed at the base into two wings running a long way down the stem, the lower ones often stalked, and six or eight inches long or more. The flowers are in a dense woolly terminal spike, a foot long or more. The corolla is yellow, slightly concave; the stamens are five in number, three hairy, the other two longer and smooth. The whole plant is very showy and attractive when growing in perfection, as it sometimes does on open moors and commons. The common name Torch-blade, or King's Taper, may have arisen from its candle-like appearance when growing by itself, pointing straightly upwards, and with its flame-like crown of flowers. Some authors account for these common names rather by the fact, that the woolly covering, which is still collected for tinder, was at one time used as wicks for tapers, especially for those employed in religious services; the very trouble entailed in collecting the material seeming to recommend it for this purpose in a church where bodily sacrifices and penances are enjoined. The Verbascum was formerly held in high repute as a medicine in disease of the lungs; it possesses some mucilaginous sedative qualities, which caused the old herbalists to believe in it. It is one of the many herbs said to stupefy or poison fish. According to an old writer, its ashes, made into soap, will restore hair which has become grey to its original colour.

RED HEMP-NETTLE.

(Fig. 69.)

GALEOPSIS LADANUM.

THIS is a pretty and not uncommon plant, belonging to the natural family Labiatae. It flowers during the whole summer and autumn in gravelly and chalky corn-fields. It is seldom above eight or nine inches high, with very spreading falling branches, covered with soft short down. The leaves are shortly stalked, narrow, ovate, or lanceolate, and very coarsely toothed. The flowers are variegated with red and yellow, sometimes nearly white; they grow in whorls around the stem, and are remarkable for the very pointed, long teeth of the calyx; which, however, are more marked in the common Galeopsis, *G. Tetrahit*.

All the British species of Hemp Nettle are pretty plants. The large flowered species, *G. versicolor*, is much more beautiful than our example, but is much less common in England. In the Highlands of Scotland it is abundant. Rare also is the downy Hemp-nettle, *G. villosa*, and is only recognized as a variety by some writers.

BLADDER-WORT

(Fig. 70.)

UTRICULARIA VULGARIS.

THE BLADDER-WORT is a pretty aquatic plant, abundant in our pools and water-channels, belonging to the family Lentibulaceæ. It is very remarkable for the buoy-like vesicles which are developed on its immersed leaves, and which serve to float the plant above the water. At certain seasons, the whole plant is submerged, and then, if we examine these vesicles we find them filled with water; gradually air becomes generated in the vesicles, expels the water, and the apertures are closed by a curious valve, so that the plant now becomes buoyant, and rises to the surface; the flowers expand, the seeds ripen, and then the living energy of the plant seems exhausted—air no longer fills the vesicles, but water takes its place, the plant sinks to the bottom, and the seeds are sown in their most fitting soil. The flowers are rather large, and of a yellow colour. The corolla has a short, conical, more or less curved spur. The leaves are pinnate with numerous segments. In the eastern counties of England the Bladder-wort is not uncommon, and in many other localities it has been found, although it must be considered as rather rare.

SEA MILKWORT

(Fig. 71.)

GLAUX MARTIMA.

SEA MILKWORT is common on nearly every sea-coast. It belongs to the family Primulaceæ, and is conspicuous from its rose-coloured tiny flowers, resting in the axils which its ovate fleshy opposite leaves form with its branched and procumbent stem. It is very pretty while growing, and is associated with some of our pleasantest sea-side rambles, and though scarcely attainable without incurring wet feet, it is worth the inconvenience ; and those who are in health must not shrink from searching the salt marshes on the coast, for many very charming botanical treasures find their homes there.

 PRIMROSE.

(Fig. 72.)

PRIMULA VULGARIS.

THE Primrose is a type of the natural order Primulaceæ. We select this plant because it is the real true primrose of our childhood—not the cowslip, with its deep yellow cups and nodding flowers—nor the oxlip, with its larger but paler blossoms ; but the sulphur-coloured primrose with which we all associate the early

days of spring, and the first ramble in the meadows or by the hedge-side. The first Primrose of the year is prized and welcomed, but when the thick tufts of its blossoms are seen, then we feel that spring is really come, and we almost fancy that no other flower ever looked so lovely or so fresh and pure. Botanically, the Primrose is an excellent example of a plant with regular monopetalous corolla, and is a good specimen for a first lesson in botany. Pretty flower as it is, all animals reject it as food excepting the pig. It seems, however, not wholly objected to by man, or woman either, for I lately saw a receipt for a primrose-pudding. A kind of wine, too, is made from the flowers, something like cowslip wine, but more delicate in flavour.

MONEYWORT, OR CREEPING LOOSE-STRIFE.

(Fig. 73.)

LYSIMACHIA NUMMULARIA.

THIS is a pretty little plant, to be seen in all hedges, or on banks, during the summer and autumn, throughout England. It belongs to the natural family Primulaceæ. The stems are prostrate, trailing to the length of one or two feet, often rooting at the nodes. The leaves are round, on very short stalks, looking almost like pieces of money; the flowers yellow, large and handsome, on short stalks—the segments of the calyx are pear-shaped. With a microscope, little pedicellate glands may be seen covering the blossoms and stamens.

73

Marywort



74

Stachel Pimpernel



75

Bog Pimpernel



76

Great Plantain



77

Stickle Spurge



78

Water Starwort



79

Black Bryon



80

Water Suet



81

Large Orchid




82

Man Orchid



83

Bee Orchid



84

Red Orchid



PIMPERNEL, OR POOR MAN'S WEATHER-
GLASS.

(Fig. 74.)

ANAGALLIS ARVENSIS.

“ Whose brilliant flower,
Closes against the approaching shower,
Warning the swain to sheltering bower,
From humid air secure.”

THIS well-known and very attractive little plant belongs to the family Primulacæ, and is a neat, much-branched, procumbent annual, six inches to near a foot long, with opposite, broadly ovate, sessile and entire leaves. The calyx divisions are pointed. The corolla is rotate, and usually of a bright red colour; sometimes it is white or pale-pink, or blue. The blue variety is as common in central and southern Europe as the red is with us, but in England it is very rare. I have, however, found it in Suffolk, in a lane near Felixstowe, where the red pimpernel grows most luxuriantly. The extreme sensitiveness of this pretty little plant to a change of atmosphere causes it to shut up its petals at the approach of rain. In fine weather it remains open from about eight in the morning till four in the afternoon. It is a common weed in the valley of the Nile, and its botanical name, which is derived from the Greek, signifies a “reviver of the spirits,” in allusion to the medical and magical properties for which it was at one time highly valued.

At present its only use seems to be as a potherb, and it is also sometimes—more especially on the Continent—eaten as a salad.

BOG PIMPERNEL.

(Fig. 75.)

ANAGALLIS TENELLA.

THIS is a rarer plant than the former species, but is equally beautiful. It is a creeping, delicate perennial plant, with very small heart-shaped or circular leaves. The flowers are very elegant, of a pale-pink colour, on long slender stalks. The corolla is very delicate in texture, and the stamens are surmounted with very woolly filaments. It flowers throughout the summer, and must be looked for in damp, boggy places, on wet mossy banks, or by the side of rivulets. It is spread over the greater part of Britain, but chiefly in the west.

PLANTAIN, OR WAY-BREAD.

(Fig. 76.)

PLANTAGO MAJOR.

THOUGH not an attractive-looking plant as we generally see it by the wayside, it is so closely associated with almost every walk along a country lane, or even high-road, that it can scarcely escape notice. The leaves are erect or spreading, broadly ovate, entire or toothed, on longish channelled stalks. The peduncles

are usually longer than the leaves, bearing a long slender spike of sessile flowers. The sepals are green in the centre, the stamens longer than the corolla; the whole flower is small, white, with pinkish anthers. In general the Plantains are regarded merely as troublesome weeds. Their leaves are eaten by some animals, but contain very little nutriment. The common name of *P. major* is undoubtedly Way-bred, not Way-bread, as it is usually spelt, from its frequency by the wayside, seeming as if bred on the road. This plant has a peculiar tendency to follow the migrations of man, as if domestically or sympathetically attached to the human race, and has followed our colonists to every part of the world; so that it has been named by the natives in some of our settlements "The Englishman's foot," for with a strange degree of certainty it is found wherever our countrymen have trod. It is a favourite food with birds, and nothing delights them more when kept in confinement than a supply of the long stalks of plantain to pick at. In the summer, when a supply of other green food can be found for our favourites, the plantain is not so much sought for; but it should be gathered and laid by in store for the winter, when it will be a treat to the little inhabitants of our aviaries. Bruised plantain-leaves are esteemed an excellent remedy for cuts and bruises, and also as an application to the bites of stinging insects. In the Highlands of Scotland an ointment is made from them, which is said to be very efficacious. There are five British species of plantain, some of which have medicinal reputation from their mucilaginous properties. *P. coronopus* has been eaten as a salad, but is too bitter to be pleasant.

PURPLE SPURGE.

(Fig. 77.)

EUPHORBIA PEPLIS.

THIS is a handsome plant belonging to the family Euphorbiaceæ. It is found only on the sea-coasts, and is becoming rare in Great Britain. It loses all its leaves before flowering, when the short main stem divides close to the base, so that the whole plant appears to consist of the forked procumbent branches lying on the sand in patches from six inches to a foot in diameter. The whole plant has a reddish or purplish hue, and that peculiar glaucous appearance so general in seashore plants. The leaves are opposite, half oblong, heart-shaped, and very thick, with small stipules at their base. The flower heads are very small. The genus *Euphorbia* is a very extensive one; there are fourteen or fifteen British species, nearly all of which have an acrid poisonous secretion. A notion prevails that milk is an antidote to this poison, but without reason, as we read in Withering's botany of a strong youth who was poisoned by drinking milk in which the plant had been boiled. As a cure for warts we can believe in the virtues of the Spurge; its astringent or acrid qualities may have some powerful effect as an external application; but old Gerarde's advice is certainly worth quoting, where he says: "These herbes by mine advise would not be received into the body, considering that there be so many good and wholeson.

potions to be made with other herbes that may be droken without perill." The ancient Britons used the Euphorbia to poison fish, which practice is continued by the Abyssinians. It is said that the fish thus killed are good for food, but this is doubtful, as there are instances of people being poisoned merely from drinking the milk of a goat which had fed on the Euphorbia.

WATER STAR-WORT.

(Fig. 78.)

CALLITRICHE VERNA.

THIS is the representative of the only genus of Water Star-worts — Callitrichaceæ — in Great Britain. This pretty plant is very suitable for the aquarium, and is found in stagnant and slowly-running water all over the country. It is easily known by its upper leaves floating on the water, and two or three pairs of them forming a little green star, hence its name. This plant forms a beautiful object under the microscope, for its leaves and stem are covered with very minute rosette-shaped bodies, which seem to supply the place of hairs in other plants. There are other species of Water Star-wort in England, but they are much more rare.

BLACK BRYONY.

(Fig. 79.)

TAMUS COMMUNIS.

THIS is an elegant climbing plant, which through the whole summer may be seen festooning our hedges and bushes. Its bright shining heart-shaped leaves clothe the trunks of many of our sturdy old trees with a verdure unequalled by any other plant. The lightness of its twining stems prevents it from injuring the branches within its grasp, for they have not the firmness or strength of the ivy, which is sometimes too close in its embrace.* The flowers are unattractive and small, but the berries of the Bryony, hanging like clusters of wild green grapes during the summer, and changing into brilliant scarlet balls in the autumn, are objects of great beauty. They are very poisonous, and must not mislead by their charming appearance. The early shoots of the plant, however, have frequently been boiled and eaten, it is said, with great relish and with no bad results. The roots have a black colour externally, hence the name of the plant. The interior is white and full of starch, which is, however, bitter

* The family Dioscoraceæ, to which the *Tamus* belongs, includes the yams; the tubers of which are well known as an article of food in South Africa and Mexico, and have lately been extensively sent to England, but, as yet, are not very popular, so great is the prejudice against any innovation on our old habits in diet.

and unwholesome. The acrid pulp of these roots has been used as a stimulating plaster.

WATER SOLDIER.

(Fig. 80.)

STRATIOTES ALOIDES.

THIS plant, so named from its sword-shaped spring leaves and fancifully military appearance, is a very ornamental aquatic plant. It belongs to the family Hydrocharideæ, and is found very abundantly in lakes and watery ditches in the east of England. In Norfolk and Suffolk it is very plentiful, and is an interesting plant to observe. It remains submerged during the greater part of the year, but raises itself to the surface on special stalks during the season for fertilizing the seeds. It forms a favourite hiding-place for multitudes of aquatic insects, as may be seen if a plant be quickly raised out of the water. The whole plant seems to resemble an aloe more than any thing else, and is equally rigid and sharp in its leaves.

EARLY ORCHIS.

(Fig. 81.)

ORCHIS MASCULA.

WE now come to a most attractive and beautiful group of wild flowers, difficult to describe, and very puzzling to the young botanist, but easily recognized

by any one who has seen and gathered them, even without knowing their scientific distinctions. The Early Orchis blossoms in the months of May and June, and may be found in any meadow or moist wood throughout Great Britain. The stem is a foot or a foot and a half high, bearing numerous showy flowers, in a loose spike, from three to six inches long, varying in colour from a pinkish purple to flesh-colour, or even white. The leaves are broad and often spotted with purple. The bracts are coloured nearly as long as the ovary, with a single rib. The upper sepal and the petals converge over the ovary, but the lateral sepals are spreading, or turned back. The lip is scarcely longer than the sepals, often slightly downy in the centre, turned back on each side with three short lobes, the middle one the largest. The tubers of all the Orchis plants contain a good deal of farinaceous nutritious matter, consisting, according to modern chemists, of a principle called Bassorin. This substance is commonly known as *saloop*, or *salep*, a word derived from the Persian name of the Orchis, which, according to Forskhale, is *sahleb*. When boiled in water, it used to be sold at the corners of the streets in London, and was a favourite drink with coalheavers, porters, and other hard-working men. It is still highly esteemed in the East; and during the Great Exhibition of 1851 was exhibited and sold as a beverage. It is said to contain more nutritious matter according to its bulk than any other vegetable substance, and that an ounce a day will sustain a man; hence it is a favourite food, from its portability, with pedestrian travellers in wild deserts and uninhabited countries.

MAN, OR MILITARY ORCHIS.

(Fig. 82.)

ORCHIS MILITARIS.

THIS is a handsome species of Orchis, from one to two feet high, with entire tubers. It is difficult to say what fancy has given rise to the name ; the red colour of the flowers or the shape of the root may have suggested a soldier-like resemblance. The leaves in the lower part of the stem vary from broadly oval to oblong ; they are usually three to five inches long. The flowers are numerous, in a dense oblong spike, with short bracts. The sepals are red or purple, and converge over the petals and column in the shape of a helmet. The lip is more or less spotted with rough, red points, and four-lobed, or rather *three-lobed*, with two entire lobes, and a third one divided in the middle into two, with a small tooth in the cleft or notch. It is found on chalky hills, on the borders of woods, and in hilly pastures. Near London it should be looked for about Dorking, Rochester, and Northfleet. There are very many other British Orchids ; all of them are pretty and attractive, and one, *A. conopsea*, has a fragrant, agreeable scent.

BEE ORCHIS.

(Fig. 83.)

OPHRYS APIFERA.

THIS plant belongs to the family Orchidaceæ, and to a section of that family most curiously eccentric in the forms of its flowers, very closely resembling the Orchies with the habits, tubers, and foliage of that group of plants ; but the flowers of *Ophrys* have no spur, and the lip is usually very curved, resembling, more or less, the body of an insect. They are often called Orchis, though not properly belonging to that genus. The Bee Orchis blossoms in July, in dry pastures, in chalky and limestone districts. In the eastern counties of England it is not uncommon. It has a brown, velvety lip, variegated with yellow, which resembles the body of a bee ; the purplish petals look like the expanded wings of an insect which had just settled on the stem of the flower. Langhorne's pretty lines have been often quoted, but are too descriptive to be omitted here :—

“See on the flow’ret’s velvet breast
 How close the busy vagrant lies,
 His thin-wrought plume, his downy breast,
 The ambrosial gold that swells his thighs.

Perhaps his fragrant load may bind
 His limbs—we’ll set the captive free;
 I sought a living bee to find,
 And found the picture of a bee.”

FLY ORCHIS.

(Fig. 84.)

OPHRYS MUSCIFERA.

THIS is a much more delicate, slender plant than the Bee Orchis, with narrow leaves and a slender spike of three or four flowers. The sepals are whitish-green. The lips brownish-purple, oblong, convex, and with pale-blue or white marks in the centre ; the two lateral lobes turned down ; the central one large, with a deep notch. The blue spot upon the base of the middle segment of the lip contributes much to the resemblance of the flower to a fly, which, indeed, is singularly close. We have also in this genus of plants the Spider Ophrys ; and there is also the Butterfly Orchis, and the Lizard Orchis ; these latter names are not, however, so well applied as those of the bee and fly.

YELLOW FLAG—WATER FLAG.

(Fig. 85.)

IRIS PSEUDACORUS.

THE FLAG plant is found in wet meadows and marshes and along water-courses throughout Europe. It belongs to the family Iridaceæ. The stem is about two feet high. The lower leaves are often much longer,

stiff and erect, and of a pale-green colour ; the upper leaves are shorter. The flowers proceed from a sheathing bract, are large, erect, and of a bright-yellow colour, two or three together. The stigmas are petal-like, rather longer than the inner segments, two-cleft at the top, with a short scale-like appendage inside at the base of the lobes. The capsule is green, from two to three inches long, with numerous pale-brown seeds. The medicinal uses of this plant are various. The juice of the fresh root is very acrid, and acts powerfully. Withering mentions a case where it was given to some swine bitten by a mad dog, and they escaped the disease ; while some others, bitten by the same dog, died with all the symptoms of hydrophobia : but I need not say that unless some evidence existed of this plant possessing active properties, Withering's cases do not prove any connection between the medicine and the cure. It is on such reasoning as this that the fallacies of Homœopathy are based. The roasted seeds are said to be a good substitute for coffee, when carefully prepared. Few plants can exceed the Iris in elegance of form ; in our gardens it is a great ornament, and by the sides of streams and lakes it should always be encouraged

“ Amid its waving swords,
In flaming gold the Iris towers.”

The Iris is undoubtedly the original of the *Fleur de lys* in the arms of France, and in many pieces of sculpture in which this device is introduced, it is not difficult to recognize it. The plant was considered in ancient times as peculiarly sacred to the Virgin Mary, as

shown in the old legend of the knight who, more devout than learned, could never retain in his memory more than two words of a prayer to the Lady Mother. These were *Ave Maria*, and with these he constantly addressed his prayer to Heaven. Night and day his prayer continued, until the good old knight died and was laid in the chapel yard of the convent, when, as a proof of the acceptance of his brief though earnest prayer, a plant of *Fleur de lys* sprang up on his grave displaying on every flower in golden letters the words *Ave Maria*. The sight induced the monks, who had despised him during his lifetime on account of his ignorance, to open his grave; and there they found the root of the plant resting on the lips of the good old soldier who lay mouldering there.

In Britain we find but two really native species of Iris, *I. pseudacorus* and *I. foetidissima*, the smell of the crushed leaves of which is thought to resemble roast beef; tastes differ, however, for Linnæus, when he gave it the specific name, must have regarded it as anything but savoury in smell. The juice is sometimes used to excite sneezing in cases of headache, but it is a very unsafe practice.

DAFFODIL—DAFFY-DOWN-DILLY.

(Fig. 86.)

NARCISSUS PSEUDONARCISSUS.

THIS is a well known pretty plant, belonging to the family Amaryllidaceæ. The name *Narcissus* seems to

be wrongly applied to this species, as it belongs, no doubt, to the *Narcissus poeticus* of the Greeks, which has a flower with a very powerful scent, a quality from which the daffodil is free. The *Narcissus* was consecrated to the Furies, who stupified their victims, hence Sophocles calls these flowers "garlands of the infernal gods." The fable of the youth *Narcissus*, after whom the plant is named, is well known to everybody; how he fell in love with his own image reflected in the water, and pined away until he was changed into the pale flower which rightfully bears his name. Our present specimen, the Daffodil, bears simply the old English name *affo dyle*, which signified "that which cometh early," and it was long before the word was corrupted into *daffodil*. It is one of our earliest spring flowers; it is rare in Scotland and Ireland, but in the south-west of England its yellow or pale lemon-coloured blossoms may be seen covering acres of land. In Cornwall they are still called Lent lilies. The root, and, to some extent, the whole plant, is poisonous, yet a useful spirit is distilled from it, which has been used as an embrocation, and also given as a medicine. Most welcome are these pretty spring flowers to us all, and in cottage gardens they add beauty and grace without expense or trouble, for they grow under almost any conditions.

"When the vales are decked with daffodils,
I hail the new reviving year,
And soothing hope my bosom fills."

WILD HYACINTH.

(Fig. 87.)

HYACINTHUS NONSCRIPTUS.

THIS well-known pretty plant is known to every child who has rambled in the fields and woods in search of wild flowers. It is sometimes called the Harebell, or Bluebell, but is quite distinct from the *Campanula rotundifolia*, the true Bluebell of Scotland. It belongs to the family Liliaceæ, and by some botanists is not called *Hyacinth* at all, but is placed with the squills, and is called *Scilla nutans*. It is the child of the spring and the denizen of the woods, whilst the *Campanula* is the pride of harvest time, and flourishes in exposed situations. The wild Hyacinth has a white bulb, full of clammy juice. The leaves are linear, shorter than the flower-stem. The stem is about a foot high, angular, with a terminal one-sided raceme of drooping blue flowers, each with a small narrow bract at the base of the pedicel. We all know the beautiful sweet-scented Hyacinth of the gardens, but nothing can be more charming than the masses of these wild blue Hyacinths sometimes seen in sheltered places, under the spreading branches of trees, forming almost a living carpeting to some shady nook. In Kew Park, acres of the green grass is changed from green to blue by the presence of these lovely blue flowers. The *Hyacinthus nonscriptus* is not the true Hyacinth of ancient Greek story; that is supposed

to have been a species of Lily, which sprung from the blood of the beautiful boy Hyacinthus, whom Apollo unfortunately killed ; so that though we cannot claim for our favourite a place in classic lore, we think that its own unassuming beauty, delicate scent, and the early associations it must recall to all hearts, will justify its introduction here. The juice contained in the root of the Hyacinth is sometimes used as starch or gum. At one time, when stiff ruffs were worn, it was much in request. It also served the purpose of the bookbinder, to fasten the covers of books securely.

GRAPE HYACINTH.

(Fig. 88.)

MUSCARI RACEMOSUM.

THE GRAPE HYACINTH belongs to the order Liliaceæ, but may be easily distinguished from our common Hyacinth, although belonging to the same family, and flowering about the same time. The leaves are narrow, lanceolate, rather thick, not stiff, from six inches to a foot long, or, when very luxuriant, $1\frac{1}{2}$ feet long. The stem is usually shorter, with a close terminal raceme, or head of small, dark-blue flowers, looking almost like little berries ; a few of the uppermost are of a paler blue, erect, much narrower, and without stamens or pistil. At one time this plant was very much cultivated in gardens. It abounds in sandy soils, and is plentiful in the eastern counties of England.

SPRING SQUILL.

(Fig. 89.)

SCILLA VERNA.

THIS plant belongs also to the family Liliaceæ, and is a pretty plant, found chiefly in the north and west coasts of England, near the sea. In Wales it has, however, been frequently met with, and on the east coast of Ireland, but rarely in Scotland. It is a delicate little plant, with a small bulb, and narrow, linear leaves, two to four inches long. The flower-stem is seldom above six inches long, with several small, erect, blue flowers, in a short terminal raceme, almost forming a corymb. The blossoms are bell-shaped, but not pendant, and of a bluish-violet colour. The foreign species of Squills have long been celebrated for their medicinal qualities, but I am not aware that any properties of a like kind have been attributed to their British relatives.

 SNAKESHEAD LILY, OR COMMON
FRITILLARY.

(Fig. 90.)

FRITILLARIA MELEAGRIS.

THIS pretty plant belongs to the Lily tribe—Liliaceæ; indeed, this is so beautiful a family of

plants, that it is difficult not to show a partiality to it when choosing wild flowers "worth notice." The Fritillary is not a common plant; it is found chiefly in the southern and eastern counties of England, and cannot be said to be wild at all in Scotland. The unexpanded blossom, somewhat fancifully, is said to resemble a snake's head; hence its common name. It has a stem about a foot high, with its leaves alternate and linear lanceolate; the single drooping flower, of a dull-red colour, marked curiously with pink and dark-purple; hence the name *Fritillaria*, from *Fritillus*, a dice-box—the frequent companion of a cheque or checkered board. It blossoms in April, in meadows and pastures throughout England, and is the only British species of the genus. Many handsome foreign species are cultivated in our gardens. The handsome Crown Imperial (*Fritillaria imperialis*) is a native of Persia, and belongs to the Tulip group of the liliaceous tribe of plants.

TURK'S-CAP LILY.

(Fig. 91.)

LILIUM MARTAGON.

THIS belongs to the Tulip group of the Lily family of plants. It is not a common plant in its wild state, though in gardens it is cultivated to great perfection and beauty. It is to be met with, however, on chalk hills and in woody places in the southern parts of England. The scent of the Lily is so powerful as

to be very annoying to some people, though in this species, and in their wild state, it is not so perceptible. The flower may readily be recognized, from its common name. The petals are reflexed, and turn over, forming a sort of turban, while the stamens appear like a tuft of feathers at the top.

MEADOW SAFFRON.

(Fig. 92.)

COLCHICUM AUTUMNALE.

IN the autumn of the year, when all nature seems losing her brilliancy, and the "sear and yellow leaf" proclaims the approach of winter, this pretty flower forms a gay carpet in the fields and meadows. It may easily be mistaken for a crocus, which, however, belongs to a different family, and blossoms in the spring of the year.

"The Crocus blows before the shrine,
At vernal dawn of St. Valentine."

The *Colchicum* belongs to the family *Melanthaceæ*, while the *Crocus* is placed with the *Iridaceæ*. The flowers of the *Colchicum* are large, of a pale-purple colour, and spring up without leaves, forcing their way through the soil, and expanding just above the ground, leaving the tubular part with the ovary and filaments enveloped in membranous sheathing spathes below the soil. Each stalk produces six or eight of these flowers.

The stamens are six, the ovaries three, each with a long thread-shaped style, and not adhering in any manner to the flower. These are succeeded by the fruit in the form of three little follicles, which slightly adhere together by their inner edge, and in the spring are elevated above the soil by their lengthened footstalk. At this time too the foliage makes its appearance in the form of an erect tuft of oblong shining sheathing leaves. Each follicle contains several oblong seeds. So like is the whole plant to the autumn Crocus, that inexperienced observers may readily mistake one for the other. They may be distinguished, however, by remembering that the Crocus has only three stamens, one style, and an inferior ovary ; while the Colchicum has six stamens, three styles, and a superior ovary. The Crocus, too, is perfectly free from those poisonous qualities which distinguish the Colchicum, and render it so useful a plant in medicine. Many species of Colchicum are cultivated for the sake of their flowers, and are badly distinguished by botanists from our specimen, which alone is of any worth in materia medica. So virulent are the poisonous properties of the *C. autumnale*, that the fingers are sometimes benumbed in preparing it, and cattle which have been driven to eat it through hunger have frequently died. It is very important for those who are employed in collecting the plant for medicinal use, to know that the active principle it contains appears to become concentrated in different parts of the plant at different seasons of the year. In June and July the root is in perfection ; in September the flowers, and the seeds in the following spring. No vegetable poison has been more advantageously applied

in medicine than this. Sir Everard Home recommended a tincture of it as efficacious in curing the gout, and it is now even more generally prescribed in rheumatism and gout than in his time. The Hemo-dactyl of the Greeks is believed to have been a species of *Colchicum* ; it was celebrated as a remedy for gout in ancient times, and its celebrity has been again revived as an ingredient in the French remedy—" *Eau Médicinale.*"

FLOWERING RUSH.

(Fig. 93.)

BUTOMUS UMBELLATUS.

THIS is a peculiarly elegant plant, belonging to the family *Alismaceæ*. It is a rush-like plant, with three-cornered, sword-shaped leaves, and umbels of handsome rose-coloured flowers containing nine stamens, a peculiarity by which it is immediately recognized among other wild flowers. The roots are regarded in Russia as a specific in hydrophobia, but experiments made with them in this country have not confirmed the accounts given of their properties by Russian physicians, and they do not seem to offer any remedy for this terrible and incurable disease. Well may the name of this beautiful plant signify the "Pride of the Water:" it is one of the most ornamental natural adornments of our lakes and rivers, and in marshy districts, where all looks barren and desolate, there may be seen the bright-coloured flowers

of the Flowering Rush, by their beauty making "the wilderness to rejoice and blossom as the rose."

Withering quotes a couplet, which is expressive and true:—

"Her rosy umbels rears the Flowering Rush,
While with reflected charms the waters blush."

COMMON ARROW-HEAD.

(Fig. 94.)

SAGITTARIA *SAGITTIFOLIA*.

THIS plant belongs to the family Alismaceæ, but differs from some other genera of that family in having unisexual flowers. The leaves, which are arrow-shaped, rise out of the water on very long stalks, the blades six or eight inches long. The flower stem is leafless, erect, and longer than the leaves—bearing on its upper part several distant whorls of rather large white flowers with a purplish tinge at the base of the petals; but so readily do they fall off, that it is difficult to preserve them. The upper flowers are those which contain the stamens, the lower ones on shorter stalks contain the pistils. There is but one locality in Scotland recorded where this Flowering Rush is growing wild; that is somewhere near Paisley. In England and Ireland it is not uncommon, and may be seen in luxuriance on the banks of the Thames, above Putney, during the summer and autumn. The roots of this plant, as well as those of other species, contain an amylaceous

matter, which is said to form a nutritious food, and is eaten for that purpose by the Chinese and Kalmuk Tartars.

CUCKOO PINT,—LORDS AND LADIES.

(Fig. 95.)

ARUM MACULATUM.

CAN we wonder at the delight of country children with this curious plant, which seems almost to be one of those things we constantly see in nature, designed to illustrate the grotesque as well as the beautiful. Its large handsome spathe, rising up amidst the elegantly-shaped spotted leaves, forms a fitting shelter for the bright-coloured spadil or flowerstalk, the lord or lady, whichever it may be, within its protecting hood. The plant belongs to the family Araceæ, and is the only representative of its family found wild in this country. It would puzzle the beginner in botany to make out the parts of the flower corresponding to those of a Primrose, but after a little study they will be easily understood. The attraction of this curious plant does not cease after the early spring, when the green leaves have faded away, and the lords and ladies and their habitation are no more seen; the little bunch of seed-like bodies about half-way down the coloured spadil, which are in fact the pistils and seed-vessels, in the autumn of the year assume a brilliant red colour, looking like a bunch of coral, as amid the withering

grass of some hedge-side they attract the notice of the passer-by ; beware, however, of being tempted to taste them. The whole plant is acrid, pungent, and poisonous. The tubers contain a sort of farinaceous substance, which, when freed from its acrid qualities, constitutes a nutritious article of diet. Large quantities are collected in Portland Island and on the dry and sunburnt districts on the banks of the Bristol Channel, and sold under the name of Portland sago. It is largely used to adulterate arrow-root. On the Continent its economical uses appear to be generally known and appreciated. Dr. Withering quotes Wedelius for the supposition that it was on this plant, under the name of Chara, on which the soldiers of Cæsar's army subsisted when encamped at Dyrrachium. A curious belief is recorded by Gerard as coming from Aristotle, that when bears were half-starved with hybernating, and have lain in their dens forty days without any nourishment but such as they get by "sucking their paws," they were completely and suddenly restored by eating this plant. Medicinally the Arum had at one time a great reputation in common with all other plants containing acrid or poisonous principles. In rheumatism, gout, and even consumption, its virtues were vaunted, but are now happily discarded. The Arum is one of those plants which exhibits the curious and interesting fact of the vegetable evolution of heat, so evident, that for some hours after the opening of the spathe it may be felt with the hand or tested with the thermometer.

SWEET-FLAG—SWEET-SEDGE.

(Fig. 96.)

ACORUS CALAMUS.

IT is, perhaps, well to conclude our selection of Wild Flowers with one which is worth notice for qualities that do not directly address the eye. Beauty of form and colour are not alone the attractions of the vegetable kingdom, and, as with many other good things in life, we do not become aware of their excellence on a superficial acquaintance. The charming qualities of this our last specimen are undiscovered until we seek for them; not in fact until we have gathered it, and bruised it in our fingers. Then comes forth the pleasant fragrance which gives it its special character. It is a reed-like plant with a creeping horizontal stem. From this springs many deep-green sword-shaped leaves about three feet long. In the midst of all is a leaf-like stem, from below the point of which comes a cylindrical or rather conical spadix of greenish flowers, which are so densely packed together, that the stalk is quite hidden. It grows abundantly in the Eastern counties of England, on the banks of rivers, and was used formerly for strewing the churches on festival days; the custom is, I believe, still preserved in the cathedral at Norwich on certain occasions. The very aromatic smell of this plant is singular amongst our native species. This property has, as might be expected, been made available in medicine. The roots

have a strong aromatic smell and a warm pungent taste, consequently they have been prescribed in many ailments where slight stimulants seemed necessary. They constitute the *Calamus aromaticus* of the shops.

Having now gone over the allotted collection of Wild Flowers, endeavouring to chronicle the chief attractions and virtues of each ; I can but feel how little has been said when compared with all that remains unsaid, but felt. I have hesitated in making these little sketches many a time, because I had not the courage to pass by others rudely, whose claims to notice seemed as great as any I have chosen. The whole field of Nature lies before us ; I can but hope that if there have been ninety-six interesting thoughts suggested by these our selected examples, they may but be inducements to still further study of the forms and functions of those beautiful creations on which a kind Providence has made so much of our welfare and happiness to depend.—“Consider the lilies of the field, how they grow ; they toil not, neither do they spin : and yet I say unto you, that even Solomon in all his glory was not arrayed like one of these.”

SYSTEMATIC CHAPTER OF CONTENTS.

FLOWERING PLANTS.

CLASS I

EXOGENS OR DICOTYLEDONS.

Subdivision I.—THALAMIFLORÆ.

	PAGE.		PAGE.
ORDER 1.—RANUNCULACEÆ.		ORDER 7.—DROSERACEÆ.	
Ranunculus bulbosus	15	Drosera rotundifolia	35
Clematis Vitalba	16	Parnassia palustris.....	44
Anemone nemorosa	17		
Adonis autumnalis	18	ORDER 8.—CARYOPHYLLACEÆ.	
Trollius Europæus	19	Dianthus Caryophyllus	37
Aquilegia vulgaris	20	Lychnis diurna	38
Aconitum Napellus	21		
ORDER 2.—NYMPHÆACEÆ.		ORDER 9.—LINACEÆ.	
Nymphæa alba	22	Linum usitatissimum.....	38
ORDER 3.—PAPAVERACEÆ.			
Papaver Argemone	25	ORDER 10.—MALVACEÆ.	
Glaucium luteum	27	Lavatera arborea	40
ORDER 4.—CRUCIFERÆ.		Malva moschata	41
Isatis tinctoria	28	Althæa officinalis	42
Draba verna	30		
Nasturtium officinale	30	ORDER 11.—HYPERICACEÆ.	
ORDER 5.—CISTACEÆ.		Hypericum calycinum	43
Helianthemum vulgare	33		
ORDER 6.—VIOLACEÆ.		ORDER 12.—GERANIACEÆ.	
Viola odorata	34	Geranium sanguineum	45
		Oxalis Acetosella	47

Subdivision II.—CALYCIFLORÆ.

PAGE.	PAGE.
ORDER 13.—LEGUMINOSÆ.	ORDER 18.—CRASSULACEÆ.
<i>Ononis spinosa</i> 48	<i>Sempervivum Tectorum</i> 57
<i>Trifolium fragiferum</i> 49	
<i>Lathyrus aphaca</i> 50	ORDER 19.—SAXIFRAGACEÆ.
ORDER 14.—ROSACEÆ.	<i>Saxifraga Hirculus</i> 58
<i>Potentilla reptans</i> 51	<i>Chrysosplenium alternifolium</i> 60
ORDER 15.—ONAGRACEÆ.	
<i>Epilobium angustifolium</i> 52	ORDER 20.—UMBELLIFERÆ.
<i>Ænothera biennis</i> 53	<i>Sium Angustifolium</i> 59
ORDER 16.—LYTHRACEÆ.	<i>Hydrocotyle vulgaris</i> 60
<i>Lythrum Salicaria</i> 55	<i>Eryngium maritimum</i> 61
ORDER 17.—CUCURBITACEÆ.	<i>Crithmum maritimum</i> 62
<i>Bryonia dioica</i> 56	<i>Pastinaca sativa</i> 64

Subdivision III.—COROLLIFLORÆ.

ORDER 21.—LORANTHACEÆ.	ORDER 28.—AQUIFOLACEÆ.
<i>Viscum album</i> 65	<i>Ilex aquifolium</i> 83
ORDER 22.—CAPRIFOLIACEÆ.	ORDER 29.—GENTIANACEÆ.
<i>Sambucus nigra</i> 69	<i>Gentiana verna</i> 85
<i>Lonicera Periclymenum</i> 70	<i>Menyanthes trifoliata</i> 86
ORDER 23.—GALIACEÆ.	ORDER 30.—CONVOLVULACEÆ.
<i>Galium verum</i> 72	<i>Convolvulus sepium</i> 87
ORDER 24.—DIPSACEÆ.	<i>Convolvulus Soldanella</i> 88
<i>Dipsacus Fullonum</i> 73	ORDER 31.—BORAGINACEÆ.
ORDER 25.—COMPOSITÆ.	<i>Echium vulgare</i> 89
<i>Cichorium Intybus</i> 74	<i>Myosotis palustris</i> 90
<i>Aster Tripolium</i> 75	ORDER 32.—SOLANACEÆ.
<i>Inula Helenium</i> 76	<i>Hyoscyamus niger</i> 93
<i>Athemis nobilis</i> 77	<i>Atropa Belladonna</i> 95
ORDER 26.—CAMPANULACEÆ.	<i>Solanum Dulcamara</i> 97
<i>Campanula rotundifolia</i> 77	ORDER 33.—SCROPHULARIA-
ORDER 27.—ERICACEÆ.	CEÆ.
<i>Vaccinium Myrtillus</i> 79	<i>Veronica Chamædrys</i> 98
<i>Erica Tetralix</i> 80	<i>Antirrhinum majus</i> 100
<i>Calluna vulgaris</i> 81	<i>Verbascum Thapsus</i> 101

PAGE.		PAGE.	
	ORDER 34.—LAMIACEÆ.		Primula vulgaris 105
Galeopsis Ladanum	103		Lysimachia nummularia 106
	ORDER 35.—LENTIBULARIÆ.		Anagallis arvensis 107
Utricularia vulgaris	104		Anagallis Tenella 108
	ORDER 36.—PRIMULACEÆ.		ORDER 32.—PLANTAGINACEÆ.
Glaux maritima	105		Plantago major 108

Subdivision IV.—MONOCHLAMYDEÆ.

ORDER 33.—EUPHORBIACÆ.	ORDER 34.—CALLITRICHACEÆ		
Euphorbia Peplis	110	Callitriche verna	111

CLASS II.

ENDOGENS OR MONOCOTYLEDONS.

ORDER 35.—DIOSCOREACEÆ.	ORDER 40.—LILIACEÆ.		
Tamus communis	112	Hyacinthus nonscriptus	121
	ORDER 36.—HYDROCHARI-	Muscari racemosum	122
	DACEÆ.	Scilla verna	123
Stratiotes Aloides	113	Fritillaria meleagris	123
	ORDER 37.—ORCHIDACEÆ.	Lilium Martagon.....	124
Orchis mascula	113		ORDER 41.—MELANTHACEÆ.
Orchis militaris	115	Colchicum autumnale	125
Ophrys apifera	116		ORDER 42.—ALISMACEÆ.
Ophrys muscifera	117	Butomus umbellatus	127
	ORDER 38.—IRIDACEÆ.	Sagittaria sagittifolia	128
Iris Pseudacorus	117		ORDER 43.—ARACEÆ.
	ORDER 39.—AMARYLLIDACEÆ.	Arum maculatum	129
Narcissus Pseudonarcissus ..	119	Acorus Calamus	131

INDEX.

	<i>Fig.</i>	<i>Page</i>		<i>Fig.</i>	<i>Page</i>
Aconitum Napellus	7	21	Convolvulus sepium	59	87
Acorus Calamus	96	131	Convolvulus Soldanella ..	60	88
Adonis autumnalis	4	18	Creeping Cinquefoil	30	51
Althæa officinalis	22	42	Crithmum maritimum....	41	62
Anagallis arvensis	74	107	Crowfoot, Bulbous	1	15
Anagallis Tenella	75	108	Cuckoo-pint	95	129
Anemone nemorosa.....	3	17			
Anemone, Wood	3	17	Daffodil or Daffy-down-		
Anthemis nobilis	51	77	dilly	86	119
Antirrhinum majus	67	100	Deadly Nightshade	64	95
Aquilegia vulgaris	6	20	Dianthus Caryophyllus ..	17	37
Arrowhead	94	128	Dipsacus Fullonum	47	73
Arum maculatum.....	95	129	Draba verna	12	30
Aster, Sea	49	75	Drosera rotundifolia	16	35
Aster Tripolium	49	75	Dwale	64	95
Atropa Belladonna	94	95	Dyer's Woad	11	28
Bedstraw, Yellow.....	46	72	Echium vulgare	61	89
Bindweed, Great	59	87	Elder.....	44	69
Bindweed, Sea	60	88	Elecampane	50	76
Bitter-sweet	65	97	Epilobium angustifolium .	31	52
Bladder-wort	70	104	Erica tetralix	54	80
Blood Geranium	25	45	Eryngium maritimum....	40	61
Bryony, Black	79	112	Euphorbia Peplis	77	110
Bryonia dioica	34	56	Evening Primrose.....	32	53
Buckbean.....	58	86			
Bugloss, Viper's	61	89	Flag, Sweet.....	96	131
Butomus umbellatus	93	127	Flag, Yellow or Water ..	85	117
Buttercup	1	15	Flax	19	38
			Forget-me-not	62	90
Callitriche verna	78	111	Fritillaria	90	123
Calluna vulgaris	55	81	Fritillaria Meleagris	90	123
Campanula rotundifolia ..	52	77	Fuller's Teasle	47	73
Campion, Red	18	38			
Carnation or Clove-pink..	17	37	Galeopsis Ladanum	69	103
Chamomile	51	77	Galium verum	46	72
Chicory.....	48	74	Gentian, Spring.....	57	85
Chrysplenium alternifo-			Gentiana verna	57	85
lium	38	60	Geranium sanguineum ..	25	45
Cichorium Intybus	48	74	Germander Speedwell	66	98
Clematis Vitalba	2	16	Glaucium luteum	10	27
Clove-pink or Carnation..	17	37	Glaux maritima.....	71	105
Clover, Strawberry	28	49	Globe Flower.....	5	19
Codlins and Cream	31	52	Grass of Parnassus	24	44
Colicum autumnale	92	125	Great Bindweed	59	87
Columbine	6	20			

	<i>Fig. Page</i>		<i>Fig. Page</i>
Harebell	52 77	Ononis spinosa	27 48
Heath	54 80	Ophrys apifera	83 116
Heather	55 81	Ophrys muscifera	84 117
Helianthemum vulgare ..	14 33	Orchis, Bee	83 116
Hemp Nettle, Red	69 103	Orchis, Early	81 113
Henbane	63 93	Orchis, Fly	84 117
Hogsbean	63 93	Orchis, Man or Military ..	82 115
Holly	56 83	Orchis macula	81 113
Holly, Sea	40 61	Oxalis acetosella	26 47
Honeysuckle	45 70		
Houseleek	35 57	Papaver Argemone	9 25
Hyacinth, Grape	88 122	Parnassia palustris	24 44
Hyacinthus nonscriptus ..	87 121	Parsnip, Wild	42 64
Hyacinth, Wild	87 121	Pastinaca sativa	42 64
Hydrocotyle vulgaris	39 60	Pennywort, Marsh	39 60
Hyoscyamus niger	63 93	Pheasant's Eye	4 18
Hypericum calycinum....	23 43	Pimpernel	74 107
Hypericum, Large-flowered	23 43	Pimpernel, Bog	75 108
		Plantago major	76 108
Ilex Aquifolium.....	56 83	Plantain	76 108
Inula Helenium.....	50 76	Poor Man's weather-glass	74 107
Iris pseudacorus	85 117	Poppy, Prickly	9 25
Isatis tinctoria	11 28	Poppy, Yellow Horn.....	10 27
		Potentilla reptans	30 51
Lathyrus aphaca	29 50	Prickly Poppy	9 25
Lavatera arborea	20 40	Primrose	72 105
Lavatera, Sea.....	20 40	Primula vulgaris	72 105
Lily, Snakeshead	90 123	Purple Loose-strife	33 55
Lily, Turk's-cap	91 124	Purple Spurge.....	77 110
Lily, White Water.....	8 22		
Lilium Martagon	91 124	Ranunculus bulbosus	1 15
Linum usitatissimum	19 38	Red Bryony	34 56
Lonicera Periclymenum..	45 70	Red Crane's-bill	25 45
Loose-strife, Creeping....	73 106	Rest Harrow	27 48
Lords and Ladies	95 129	Rock Cist	14 33
Lysimachia Nummularia .	73 106	Rock Rose	14 33
Lythrum Salicaria	33 55	Rush, Flowering	93 127
Mallow, Marsh	22 42	Saffron, Meadow	92 125
Mallow, Musk	21 41	Sagittaria sagittifolia	94 128
Mallow, Tree	20 40	St. John's Wort	23 43
Malva moschata	21 41	Sambucus nigra	44 69
Marsh Pennywort.....	39 60	Samphire	41 62
Marsh Saxifrage	36 58	Saxifraga Hirculus	36 58
Marsh Trefoil	58 86	Saxifrage, Golden.....	38 60
Meadow Saffron	92 125	Saxifrage, Marsh	36 58
Menyanthes trifoliata....	58 86	Scilla verna	89 123
Mistletoe	43 65	Sea Aster	49 75
Monkshood	7 21	Sea Milkwort	71 105
Moneywort.....	73 106	Sempervivum Tectorum ..	35 57
Mullein, Great	68 101	Sium angustifolium	37 59
Muscari racemosum	88 122	Snake's-head Lily.....	90 123
Myosotis palustris	62 90	Snapdragon	67 100
		Solanum Dulcamara	65 97
Narcissus pseudonarcissus	86 119	Spurge, Purple	77 110
Nasturtium officinale	13 30	Squill, Spring	89 123
Nymphaea alba	8 22	Starwort, Water	78 111
		Stratiotes aloides	80 113
Oenothera biennis	32 53	Strawberry Clover	28 40
Old Man's Beard	2 16		

	<i>Fig.</i>	<i>Page</i>		<i>Fig.</i>	<i>Page</i>
Sundew	16	35	Water Lily, White.....	8	22
Sweet Sedge	96	131	Water Parsnip	37	59
Tamus communis	79	112	Water Soldier.....	80	113
Teazle, Fuller's	47	73	Water Starwort.....	78	111
Traveller's Joy	2	16	White Water Lily	8	22
Tree Mallow	20	40	Whitlow Grass	12	30
Trifolium fragiferum	28	49	Whortleberry, Black	53	79
Trollius Europæus	5	19	Wild Bryony	34	56
Utricularia vulgaris	70	104	Wild Parsnip	42	64
Vaccinium Myrtillus	53	79	Willow Herb	31	52
Verbascum Thapsus.....	68	101	Wind Flower	3	17
Veronica Chamædrys	66	98	Woad, Dyer's	11	28
Viola odorata	15	34	Wolf's-bane	7	21
Violet, Sweet	15	34	Wood Anemone.....	3	17
Viper's Bugloss	61	89	Woodbine	45	70
Virgin's Bower	2	16	Wood-sorrel	26	47
Viscum Album	43	65	Woody Nightshade	65	97
Water-cress	13	30	Yellow Bedstraw	46	72
			Yellow Vetchling, or Yel- low Pea	29	56

THE END

Small 8vo., nearly 300 cuts, price 4s., cloth.

Half-hours in the Green Lanes.

A Book for a Country Stroll. By J. E. TAYLOR, F.L.S., F.G.S.

CONTENTS.

By a Tarnside.	Beetles and other Insects in the Green Lanes.
Fishes, Mollusca, and other Objects in the Tarn.	Flowering Plants in the Green Lanes.
Reptiles in the Green Lanes.	Grasses in the Green Lanes.
Birds in the Green Lanes.	Ferns in the Green Lanes.
Mammals in the Green Lanes.	Mosses in the Green Lanes.
Butterflies and Moths in the Green Lanes.	Fungi in the Green Lanes.
Shells and Slugs in the Green Lanes.	Lichea in the Green Lanes.

By the same Author.

Fcap., 8vo., Fully Illustrated, price 4s.

Half-hours at the Seaside.

CONTENTS.

Half-an-hour with the Waves.	Preparations	Half-an-hour with the Sea-Anemones.
” ”	Sea-Weeds.	” ” Sea-Mats and Squirts.
” ”	Sponges.	” ” Sea Urchins and Starfish.
” ”	Sea Worms.	” ” Shell-fish.
” ”	Corallines.	” ” Crustacea.
” ”	Jelly-Fish.	

By the same Author.

Now ready, Fcap., 8vo., Fully Illustrated, price 4s.

Geological Stories.

A Series of Autobiographies in Chronological order, being the Stories of

A Piece of Granite.	A Piece of Purbeck Marble.
” ” Quartz.	” ” Chalk.
” ” Slate.	A Lump of Clay.
” ” Limestone.	A Piece of Lignite.
” ” Sandstone.	The Crags.
” ” Coal.	A Boulder.
” ” Rock-Salt.	A Gravel-Pit.
” ” Jet.	

“This is a most amusing work, exceedingly well illustrated.”—*Evening Standard*.
 “Every page teems with instruction. We warmly commend the book, to the young especially.”—*Suffolk Chronicle*.

Fcap. 8vo., price 2s. 6d. plain, 4s. coloured.

Half-hours with the Microscope.

By EDWIN LANKESTER, M.D. Illustrated by 250 Drawings from Nature by TUFFEN WEST. New Edition, much enlarged with full Description of the various parts of the Instrument, Chapter on the Polariscope, and Coloured Frontispiece.

Fcap. 8vo., cloth, with Illustrations on Stone and Wood, price 2s. 6d.

Half-hours with the Telescope.

Being a popular Guide to the Use of the Telescope as a means of Amusement and Instruction adapted to inexpensive Instruments. By RICHARD A. PROCTOR, B.A., F.R.A.S.

Demy 4to., cloth, price 5s.

Half-hours with the Stars.

A Plain and Easy Guide to the Knowledge of the Constellations. Showing in Twelve Maps the Position of the principal Star-Groups night after night throughout the Year. With Introduction, and a separate Explanation of each Map. True for every year. By R. A. PROCTOR, B.A., F.R.A.S., Author of "Half-hours with the Telescope."

The only series from which the elevation and bearing of a star, at any hour of any night, can be found at once.

"Mr. Proctor has done a great deal already to popularise astronomy by his writings, as he has also achieved much by his original investigations to advance our knowledge of the heavenly bodies."—*Popular Science Review*.

"Mr. Proctor combines the rare qualities of a profound investigator, a good scholar, and a patient master; and we think he has done more to render astronomy popular than any other writer of the present day."—*Scientific Opinion*.

TO NATURALISTS, MICROSCOPISTS, AND FIELD CLUBS.

Fcap. 8vo., cloth, 114 Illustrations, price 2s. 6d.


The Collector's Handy-Book of Algæ.

DIATOMS, DESMIDS, FUNGI, LICHENS, MOSSES, &c.; with Instructions for their Preparation and the Formation of an Herbarium. By JOHANN NAVE. Translated and Edited by the Rev. W. W. SPICER, M.A.

Fully Illustrated, price 4s. coloured by hand; 2s. 6d. plain.

The British Ferns.

(A Plain and Easy Account of.) Together with their Classification, Arrangement of Genera, Structure, and Functions, directions for out-door and in-door Cultivation, &c. By Mrs. LANKESTER.



Boston Public Library
Central Library, Copley Square

Division of
Reference and Research Services

The Date Due Card in the pocket indicates the date on or before which this book should be returned to the Library.

Please do not remove cards from this pocket.

B.P.L. Bindery
JUL 24 1879

